

Minima in branching random walks

L. Addario-Berry, B.A. Reed

July 10, 2008

Abstract

Given a branching random walk, let M_n be the minimum position of any member of the n 'th generation. We calculate $\mathbf{E}M_n$ to within $O(1)$ and prove exponential tail bounds for $\mathbf{P}\{|M_n - \mathbf{E}M_n| > x\}$, under quite general conditions on the branching random walk. In particular, together with work of Bramson (1978b), our results fully characterize the possible behavior of $\mathbf{E}M_n$ when the branching random walk has bounded branching and step size.

1 Introduction

The object of study in this paper is a supercritical branching random walk (or *tree-indexed random walk*) – which we view as a Galton-Watson tree T with root r and offspring distribution B , each of whose edges e has been augmented (or *labelled*) with an independent copy X_e of a random variable X (which we call the *step size*). For the formal details of a probabilistic construction of branching random walks, see, e.g., Harris (1963). When X is non-negative, this is often called an age-dependant branching process.

We assign to each node v of T the “displacement” (or *label*) S_v , which is the sum of the edge labels on the path from r to v . The *depth* of a node $v \in T$ is the number of edges on the path from r to v . One of the most well-studied parameters associated with branching random walks is the *minimum after n steps*, i.e., the minimum value of S_v over all nodes v having depth n in T ; we denote this quantity M_n . When X is non-negative, M_n may be viewed as the time at which the first element of the n 'th generation is born. For several special choices of branching random walk, M_n also turns out to be closely linked to the performance of data structures that arise in computer science; this connection is explored in depth by Devroye (1998).

The starting point for our research is the Biggins-Hammersley-Kingman theorem, which provides a law of large numbers for the minimum displacement M_n of a branching random walk after n steps. (We set $M_n = \infty$ if the process does not survive for n generations.) In a sequence of papers, each building on the results of the last, Hammersley (1974), Kingman (1975) and Biggins (1976) showed that under suitable

conditions on the exponential moment of X , there is a finite constant γ such that, conditional on the survival of the branching process,

$$M_n/n \rightarrow \gamma \quad \text{almost surely.}$$

When Hammersley initiated this research into the first-order behavior of M_n , he posed several questions to which complete answers remain unknown. In particular, he asked when more detailed information about $M_n - \gamma n$ than that given by the above law of large numbers can be found, about the expectation of M_n , and about whether the higher centralized moments of M_n (and particularly the variance) are bounded. Our goal in this paper is to provide answers to Hammersley's questions for a broad range of branching random walks. Before we state our results, however, we briefly discuss some previous work on this and related problems.

In a remarkable paper, Bramson (1978a) derived extremely precise information about the maximum displacement of *branching Brownian motion*. In this model, an initial particle starts at position 0 on the real line and begins a standard Brownian motion with variance 1. The particle decays according to an exponential mean 1 clock; when the clock goes off, the particle splits into two, each of which continues an independent Brownian motion and each of which independently decays according to an exponential mean 1 clock. This process is continued forever. Bramson studied the maximum displacement M_t^{br} of any particle after time t (of course, his results apply immediately to the minimum, by symmetry). He showed that the median m_t^{br} of M_t^{br} satisfies

$$m_t^{\text{br}} = \sqrt{2}t - \frac{3 \log t}{2} + c + o(1), \quad (1)$$

where c is a fixed constant, and additionally showed that $M_t^{\text{br}} - m_t^{\text{br}}$ converges in distribution. (In this paper, by \log we always mean the natural logarithm.) It is worth noting that Lifshits (2007) has apparently (see Hu and Shi (2007)) provided an example of a branching random walk for which M_n minus its median is tight but does not converge in distribution, so in general a result as strong as Bramson's will not hold in the branching random walk setting.

Since Bramson's work, it has been expected that for at least some branching random walks, the median m_n of M_n should exhibit similar behavior. However, such results have been slow in coming. McDiarmid (1995) proved (among other results) that for a wide class of branching random walks, $M_n - \gamma n = O(\log n)$ almost surely. Bachmann (1998, 2000) studied the *tightness* of the family $\{M_n - m_n : n \in \mathbb{N}\}$, where m_n is the median of M_n , showing that the aforementioned family is tight when X has logarithmically concave distribution function. (We have not stated all the conditions on X and B required for his result.) More recently, Bramson and Zeitouni (2006, 2007) also proved tightness of the family $\{M_n - m_n : n \in \mathbb{N}\}$, under a strong local condition on the lower tail of X .

In the course of studying the height of random binary search trees (RBSTs), Reed (2003) proved that for a branching random walk with deterministic binary branching and exponential mean 1 step size, there exist explicit constants $\alpha > 0$, $\beta > 0$ such that

$$m_n = \alpha n + \beta \log n + O(1), \quad (2)$$

the first such result for any branching random walk; Reed additionally proved exponential tail bounds for $\mathbf{P}\{|M_n - m_n| > x\}$. We remark that the difference in sign of the

logarithmic terms in (1) and (2) is because the former equation bounds a maximum, whereas the latter equation bounds a minimum. Drmota (2003) has also published an alternative proof of Reed’s result, using completely different techniques. Chauvin and Drmota (2006) have since extended Drmota’s techniques to show exponential tail bounds for M_n around m_n when (among other conditions) the step size X satisfies what they call an *intersection property*. In particular, their result implies that such tail bounds hold when X has logarithmically concave distribution function. In a series of papers, Broutin and Devroye (2006); Broutin et al. (2008) have recently extended the Hammersley-Kingman-Biggins theorem to a wide range of branching random walks in which dependency is allowed among the steps and in the “cutoff” at which the minimum is measured is not deterministically n , but is determined by a second family of random variables coexisting on the edges of the Galton-Watson tree T .

The results and techniques of Bramson (1978a) and Reed (2003), in particular, serve as guidance and inspiration for the work of this paper. For our approach, the asymptotic behavior of M_n and of its median m_n are best characterized in terms of the behavior of the *logarithmic moment generating function* (LMGF) Λ . For a real random variable X , we define the LMGF $\Lambda = \Lambda_X$ of X by

$$\Lambda(t) = \Lambda_X(t) = \log \mathbf{E} \{ e^{tX} \}.$$

To better understand the utility of the function Λ in studying M_n , we first recall Chernoff’s bounding technique: if $S_n = \sum_{i=1}^n X_i$ is a sum of n independent copies of X , then for any $c \in \mathbb{R}$ and $u > 0$, by using Markov’s inequality we have

$$\mathbf{P} \{ S_n \leq cn \} = \mathbf{P} \{ e^{-uS_n} > e^{-ucn} \} \leq \frac{\mathbf{E} \{ e^{-uS_n} \}}{e^{-ucn}} = \left[\mathbf{E} \{ e^{-u(X-c)} \} \right]^n,$$

so

$$\mathbf{P} \{ S_n \leq cn \} \leq \left[\inf_{u>0} \mathbf{E} \{ e^{u(X-c)} \} \right]^n. \quad (3)$$

If there is u such that $\mathbf{E} \{ e^{u(X-c)} \} < (\mathbf{E}B)^{-1}$ (where B is the branching distribution) then it follows that $\mathbf{P} \{ S_n \leq cn \} \leq (\mathbf{E}B)^{-n}$, and it is fairly easy to show, using this fact and straightforward bounds on the growth of T , that $\liminf_{n \rightarrow \infty} M_n/n > c$. (To show that in this case $\lim_{n \rightarrow \infty} M_n/n$ actually exists and to determine the limit takes more work.)

The optimal choice for u in (3) is that for which $\Lambda'(u) = c$, as may be informally seen by differentiating $\mathbf{E} \{ e^{u(X-c)} \}$ with respect to u . Choosing u in this fashion, and writing $\Lambda'(u)$ in place of c , gives the bound

$$\mathbf{P} \{ S_n \leq \Lambda'(u)n \} \leq \left[\mathbf{E} \{ e^{uX} \} \cdot e^{-u\Lambda'(u)} \right]^n = e^{-n(u\Lambda'(u) - \Lambda(u))}. \quad (4)$$

It turns out that the upper bound given by (4) is almost tight; this is the substance of the “exact asymptotics for large deviations” first proved by Badahur and Rao (1960), and is the reason that the behavior of Λ is key to our investigation. As our proof leans strongly upon the exact asymptotics for large deviations, we now take the time to formally introduce this result.

Given X , S_n and Λ as above, let D_Λ be the set of values t for which $\Lambda(t)$ is finite, and let D_Λ° be the interior of D_Λ . We say X is a *lattice random variable* with period $p > 0$ if there is a constant x such that $(X - x)/p$ is almost surely integer-valued, and p is the largest real number for which this holds. We now state the asymptotic estimates for large deviations as they appear in Dembo and Zeitouni (1992) (as Theorem 3.7.4):

Theorem 1 (Badahur and Rao (1960)). *Let $S = \{S_n\}_{n \in \mathbb{N}}$ be a random walk with step size X , and define Λ and D_Λ^0 as above. Choose any $t \in D_\Lambda^0$ with $t < 0$ and any function $g(n)$ tending to plus infinity with n ; then if X is non-lattice then for any $a \in \mathbb{R}$ with $a \leq \sqrt{n}/g(n)$,*

$$\mathbf{P} \{S_n \leq \Lambda'(t)n - a\} = (1 + o(1)) \cdot \frac{e^{at - n(t\Lambda'(t) - \Lambda(t))}}{\sqrt{\Lambda''(t)} \cdot 2\pi n}, \quad (5)$$

uniformly over a in the above range. Furthermore, if X is a lattice random variable with period p , then for any a which is a multiple of $1/p$ and for which $a \leq \sqrt{n}/g(n)$, the same result holds.

In fact, this theorem is stated with a constant in (Dembo and Zeitouni, 1992), but the proof in that book yields without modification the above formulation. Theorem 1 contains two cases: the values of a for which (5) holds depend on whether X is lattice or non-lattice. Technically, this fact necessitates a case analysis in our proof, but as the two cases are virtually identical, we do not bother with both of them. *For the remainder of the paper, we assume that the step size X is non-lattice.*

We will shortly have use of the following corollary of Theorem 1. Let $f(t) = t\Lambda'(t) - \Lambda(t)$.

Corollary 2. *For $0 \geq t \in D_\Lambda^0$, $f(t)$ is strictly decreasing. Furthermore, if X is bounded from below then $\lim_{t \rightarrow -\infty} f(t) = \log(1/\mathbf{P}\{X = \text{ess inf } X\})$ (where we interpret the right-hand-side as ∞ if $\mathbf{P}\{X = \text{ess inf } X\} = 0$).*

Proof. The fact that $f(t)$ is decreasing when $t \leq 0$ follows from Dembo and Zeitouni (1992), Lemma 2.2.5 and Exercise 2.2.24. To prove the second assertion, first assume without loss of generality that $\text{ess inf } X = 0$. It is then an easy exercise to see that $\lim_{t \rightarrow -\infty} \Lambda'(t) = 0$, so for any small enough $\epsilon > 0$, by choosing t large enough we may apply Theorem 1 to obtain asymptotics for $\mathbf{P}\{S_n \leq \epsilon n\}$. Since for any $\epsilon > 0$, $\mathbf{P}\{S_n \leq \epsilon n\} \geq \mathbf{P}\{X = 0\}^n$, it is then immediate from Theorem 1 that $\lim_{t \rightarrow -\infty} f(t)$ must be at most $\log(1/\mathbf{P}\{X = 0\})$.

To see that in fact equality holds, choose any small $\epsilon > 0$ and any $c > \epsilon$. If we are to have $S_n \geq \epsilon n$ then at most an ϵ/c proportion of the X_i can satisfy $X_i \geq c$. Letting $p = p(c) = \mathbf{P}\{X \geq c\}$, it follows that $\mathbf{P}\{S_n \geq \epsilon n\} \leq \mathbf{P}\{\text{Bin}(n, p) \geq (1 - \epsilon/c)n\}$. Since $p(c) \rightarrow \mathbf{P}\{X = 0\}$ as $c \rightarrow 0$, letting $c \rightarrow 0$ and $\epsilon \rightarrow 0$ in such a manner that $1 - \epsilon/c \rightarrow 1$, it follows from Theorem 1 and standard binomial estimates that $\lim_{t \rightarrow -\infty} f(t)$ must be at least $\log(1/\mathbf{P}\{X = 0\})$. \square

Using the above exact asymptotics, it turns out that one can prove an analogue of Bramson's result for a wide range of branching random walks. The primary result of this paper is:

Theorem 3. *Consider a supercritical branching random walk with branching distribution B and non-constant step size X , and suppose that the following conditions hold:*

- (I) *there exists an integer $d \geq 2$ such that $\mathbf{P}\{B \leq d\} = 1$;*
- (II) *there exists a real number $u > 0$ such that $\mathbf{E}e^{uX} < \infty$; and*
- (III) *there exists $t \in D_\Lambda^0$, $t < 0$ such that $t\Lambda'(t) - \Lambda(t) = \log(\mathbf{E}B)$.*

Let \mathcal{S} be the event that the branching random walk survives. Then

$$\mathbf{E}\{M_n \mid \mathcal{S}\} = \Lambda'(t)n - \frac{3}{2t} \log n + O(1), \quad (6)$$

and there exist constants $C > 0$, $\delta > 0$ depending only on X , such that for all $x \in \mathbb{R}$,

$$\mathbf{P}\{|M_n - \mathbf{E}\{M_n \mid \mathcal{S}\}| \geq x \mid \mathcal{S}\} \leq Ce^{-\delta x}. \quad (7)$$

A few remarks on the conditions appearing in Theorem 3 are in order. Condition (III) is perhaps the least intuitive. For example, if X is bounded from below, then by Corollary 2, (III) simply requires that $\mathbf{P}\{X = \text{ess inf } X\} < 1/\mathbf{E}B$. Together with a result of Bramson (1978b), this shows that condition (III) can not be removed. In particular, Bramson (1978b) has shown that if X is bounded from below and $\mathbf{P}\{X = \text{ess inf } X\} = \frac{1}{\mathbf{E}B}$, then $\mathbf{E}M_n = (\text{ess inf } X) \cdot n + O(\log \log n)$ (Dekking and Host (1991) contains further information about behavior of M_n in this case). The following theorem highlights another situation in which $\mathbf{E}M_n$ does not have the form $\alpha n + \beta \log n + O(1)$:

Theorem 4. *Given a supercritical branching random walk satisfying conditions (I) and (II), above, if X is bounded from below and $\mathbf{P}\{X = \text{ess inf } X\} > 1/\mathbf{E}B$, then*

$$\mathbf{E}\{M_n \mid \mathcal{S}\} = (\text{ess inf } X) \cdot n + O(1)$$

and there exist constants $C' > 0$, $\delta' > 0$ depending only on X such that for all $x > 0$, $\mathbf{P}\{M_n > (\text{ess inf } X) \cdot n + x \mid \mathcal{S}\} \leq C'e^{-\delta'x}$.

Theorem 4 is not particularly difficult. It may be proved using the techniques developed by Hammersley, Kingman, and Biggins in the course of proving the Hammersley-Kingman-Biggins theorem, and is also essentially contained within Dekking and Host (1991). We provide a proof that allows us to highlight, in a simplified setting, a technique we later use in proving Theorem 3. We remark that together with the aforementioned result of Bramson (1978b), Theorems 3 and 4 completely determine the possible behavior of M_n in the case that X is bounded from below.

If X is not bounded from below, then condition (III) essentially imposes that the left tail of X decays “sufficiently” (and in particular, at least exponentially) quickly. In particular, if X is not bounded from below and all exponential moments $(X - \mathbf{E}X)_-$ are finite, then (III) will hold.

Condition (II) is essentially necessary for our result to hold; in general, a relaxation of condition (II) may require a corresponding relaxation of the upper tail bound in (7). In particular, it may be the case that

$$\mathbf{P}\{\text{The root } r \text{ has exactly one child} \mid \mathcal{S}\} > 0,$$

in which case it is not hard to see that the upper tail of M_n can decay no more quickly than the upper tail of X .

Finally, condition (I) is a requirement imposed by our particular use of the second moment method in the course of the proof. Independently of the current work, Hu and Shi (2007) have shown that assuming bounded step size, assuming that (III) holds and that $\mathbf{E}\{B^{1+\epsilon}\} < \infty$ for some $\epsilon > 0$, then $(M_n - \Lambda'(n))/\log n \rightarrow (-3/2t)$ in probability on \mathcal{S} but that, surprisingly, $(M_n - \Lambda'(n))/\log n$ does *not* converge almost surely. It seems very likely, particularly in view of the former result, that condition (I) should not be necessary for the results of Theorem 3 to hold. However, we would expect different behavior in the case $\mathbf{E}B = \infty$.

1.1 Our approach

We observe that it clearly suffices to prove Theorem 3 when $\mathbf{E}X = 0$, and we shall hereafter assume this is the case. To begin to describe our approach, we observe that for $\lfloor (\mathbf{E}B)^n \rfloor$ independent random walks with step size X , the expected minimum value $\mathbf{E}M_n^{\text{ind}}$ of any node w at depth n in this “forest of random walks” is within $O(1)$ of the smallest m for which

$$\mathbf{E} \{ |\{ \text{nodes } w \text{ at depth } n \text{ such that } S_w \leq m \}| \} \geq 1.$$

In other words, $\mathbf{E}M_n^{\text{ind}}$ is within $O(1)$ of $\inf\{m : \mathbf{P}\{S_w \leq m\} \geq 1/\lfloor (\mathbf{E}B)^n \rfloor\}$, a value we can easily derive to within $O(1)$ using Theorem 1. To get an idea why this is the case, choose $t \in D_\Lambda^o$, $t < 0$ and such that $t\Lambda'(t) - \Lambda(t) = \log \mathbf{E}B$. For any real number a which is $o(\sqrt{n})$, by Theorem 1 and a union bound we have

$$\begin{aligned} \mathbf{P} \left\{ M_n^{\text{ind}} \leq \Lambda'(t)n - a \right\} &\leq (1 + o(1)) \lfloor (\mathbf{E}B)^n \rfloor \cdot \frac{e^{at - n(t\Lambda'(t) - \Lambda(t))}}{\sqrt{\Lambda''(t) \cdot 2\pi n}} \\ &= (1 + o(1)) \frac{\lfloor (\mathbf{E}B)^n \rfloor}{(\mathbf{E}B)^n} \frac{e^{at}}{\sqrt{\Lambda''(t) \cdot 2\pi n}} \\ &= e^{at - (\log n)/2 - O(1)}, \end{aligned}$$

so $\mathbf{P}\{M_n^{\text{ind}} \leq \Lambda'(t)n - a\}$ is exponentially small in $((\log n)/2t - a)$ when $a = o(\sqrt{n})$ and $a > 0$. Similarly, if $M_n^{\text{ind}} > \Lambda'(t)n - a$ then each of the $\lfloor (\mathbf{E}B)^n \rfloor$ random walks S must have $S_n \geq \Lambda'(t)n - a$. By the independence of the random walks and by Theorem 1, we have

$$\begin{aligned} \mathbf{P} \left\{ M_n^{\text{ind}} > \Lambda'(t)n - a \right\} &= (1 - \mathbf{P}\{S_n \leq \Lambda'(t)n - a\})^{\lfloor (\mathbf{E}B)^n \rfloor} \\ &= \left(1 - (1 + o(1)) \frac{e^{at - (\log n)/2 - O(1)}}{(\mathbf{E}B)^n} \right)^{\lfloor (\mathbf{E}B)^n \rfloor} \\ &\leq \exp\{-(1 + o(1))e^{at - (\log n)/2 - O(1)}\}, \end{aligned}$$

so $\mathbf{P}\{M_n^{\text{ind}} \geq \Lambda'(t)n - a\}$ is doubly exponentially small in $(a - (\log n)/2t)$ when $a = o(\sqrt{n})$ and $a < 0$. These inequalities do not quite yield a bound on $\mathbf{E}M_n^{\text{ind}}$, but they do show that an extremely high proportion of the probability mass of M_n^{ind} lies near $\Lambda'(t)n + (\log n)/2t$, and by combining these bounds with a Chernoff bound for the lower tail of M_n^{ind} , it is not hard to show that in fact

$$\mathbf{E} \left\{ M_n^{\text{ind}} \right\} = \Lambda'(t)n - \frac{\log n}{2t} + O(1).$$

We denote by $m^* = m_n^*$ the quantity $\Lambda'(t)n - \log n/2t$, and call m^* the *breakpoint*. Whenever we write m^* without subscript we always mean m_n^* . By linearity of expectation, it follows easily from Theorem 1 that in the branching random walk,

$$\begin{aligned} \mathbf{E} \{ |\{w \text{ at depth } n : S_w \leq m\}| \} &\gg 1 && \text{when } m - m^* \gg 1, \\ \mathbf{E} \{ |\{w \text{ at depth } n : S_w \leq m\}| \} &\ll 1 && \text{when } m^* - m \gg 1. \end{aligned}$$

Intuitively, then, the equation (6) can be understood by splitting the term $(3/2t) \log n$ into two pieces and writing

$$\mathbf{E} \{ M_n \mid \mathcal{S} \} = m^* - \frac{\log n}{t} + O(1); \tag{8}$$

the term $(\log n)/t$ must then be explained by the dependence in the branching random walk model. The bulk of the work of this paper is in understanding and explaining why this dependence should contribute a term of just this form.

At a high level, to explain this term, we shall apply the second moment method to bound the number of certain “special” nodes of T . We will introduce a notion of “goodness” of nodes; whether or not a node v is good will depend only on the shape of the random walk from r to v . We shall study the properties of the good subset G of the nodes of T at depth n , first showing that $|G|$ is tightly concentrated around its mean, and then, with some additional work, showing that $\mathbf{E}M_n$ is in fact within $O(1)$ of $\inf_m \mathbf{E} \{ |\{w \in G : S_w \leq m\}| \} \geq 1$. To begin to make this more concrete, we first explain the key property that “good” nodes will satisfy.

1.2 Leading nodes

Given exchangeable random variables X_1, \dots, X_n with associated partial sums S_1, \dots, S_n , we say that S_n is *leading* (or that S is leading after n steps) if

$$S_i \geq \mathbf{E} \{S_i \mid S_n\} \quad \text{for all } i = 1, \dots, n; \quad (9)$$

equivalently, if $S_i \geq S_n \cdot (i/n)$ for all $i = 1, \dots, n$. (This terminology was introduced by McDiarmid (1995).) Similarly, we say that S_n is *strictly leading* if the inequality in (9) is strict for all $i = 1, \dots, n$. Given a node v at depth n in T , we say that v is a leading (resp. strictly leading) node if the random walk from the root to v is leading (resp. strictly leading).

The intuition for why leading nodes are useful may be gleaned from examples (A) and (C) of the preceding section. If a node v at depth n is leading, then knowing that $S_v \leq m$ does not increase the *expected* number of nodes w at depth n with $S_w \leq m$ by too much. This fact allows us to use the *second moment method* (i.e., some variant of Chebyshev’s inequality) to bound from below the probability that some such node exists when the expected number of such nodes is $\Omega(1)$. (This line of argument via Chebyshev’s inequality is quite common in combinatorial settings; see, e.g., Alon and Spencer (2000), Chapter 4.)

It turns out that $\mathbf{E} \{M_n\}$ is within $O(1)$ of the smallest value m_n for which the expected number of leading vertices v at depth n with $S_v < m_n$ is at least 1. Though it may not be immediately obvious, the assertion of the previous sentence is in fact equivalent to (6); we now explain this equivalence.

To bound the probability that a vertex is leading, we use a classic combinatorial technique introduced by Andersen (1953) (and first used in the context of branching random walk by Devroye and Reed (1995)), that we call a *rotation argument*, which involves considering cyclic permutations of the random variables X_1, \dots, X_n ; we will explain this approach in more detail in Section 3.

By this method, we will straightforwardly be able to show that for all $v \in N_n$ and for values $m \leq \mathbf{E}M_n$ that are not too far from the breakpoint m^* ,

$$\mathbf{P} \{S_v \leq m \text{ and } v \text{ is a leading node}\} = \Theta \left(\frac{\mathbf{P} \{S_v \leq m\}}{n} \right). \quad (10)$$

(We shall make this more formal in Section 3.) It follows that for such m ,

$$\begin{aligned} \mathbf{E} \{|\{v \in N_n, S_v \leq m, v \text{ is leading}\}|\} &= \Theta \left(\frac{\mathbf{E} \{|\{v \in N_n, S_v \leq m\}|\}}{n} \right) \\ &= \Theta \left(\frac{(\mathbf{E}B)^n}{n} \cdot \mathbf{P} \{S_v \leq m\} \right), \end{aligned} \quad (11)$$

so we are in fact asserting that $\mathbf{E}M_n$ is within $O(1)$ of the smallest value m for which, for nodes v at depth n in T , $\mathbf{P} \{S_v \leq m\} \geq (\mathbf{E}B)^n/n$. It follows immediately by Theorem 1 that m is within $O(1)$ of $m^* + (\log n)/t$, in accordance with (6) and (8). We may then view the term $(\log n)/t$ as *the correction required in order to find a leading node*. One of the key steps in proving Theorem 3 will be to understand the shape of random walks that end in leading nodes, and in particular how much time such walks spend near their (conditional) means.

1.3 A little notation and a basic fact

Let T_n be the subtree of T consisting of all nodes of depth at most n . We may view T_n as a subtree of a rooted, labeled d -ary tree T'_n with n levels and with root r , in the following manner. For a given node v of T'_n with children w_1, \dots, w_d , let B_v be a copy of the offspring random variable B , and let $X_{v,1}, \dots, X_{v,B_v}$ be independent copies of X . Let σ_v be a uniformly random permutation of $\{1, \dots, d\}$; we assign label $X_{v,i}$ to edge $vw_{\sigma_v(i)}$ for $1 \leq i \leq B_v$ and assign label ∞ to edge $vw_{\sigma_v(i)}$ for $B_v < i \leq d$. (This permutation gives the model a useful symmetry property that will be explained shortly.) We repeat this procedure for all nodes of T'_n , and label each node v with the sum S_v of all edge labels on the path from r to v (setting $S_v = \infty$ if any of these labels are infinite). With this labeling, the subtree of T'_n induced by nodes v with $S_v < \infty$ is distributed precisely as T_n , and we hereafter view $T_n = T_n(T'_n)$ as a subtree of such a d -ary tree T'_n .

For $1 \leq i \leq n$ and for any $m \in \mathbb{R}$, let N_i (resp. N'_i) be the set of nodes of T_n (resp. T'_n) at depth i , and let $N_{i,m}$ be those nodes v in N_i with $S_v \leq m$. To each $v \in T'_n$ we assign a label S_v that is the sum of the edge labels X_e on the path from r to v – so it is possible that $S_v = \infty$ for some nodes v . We observe that for any integer $m \geq 0$, M_n is the smallest $m \leq +\infty$ for which $N_{n,m} \neq \emptyset$.

For the sake of our analysis, it will be useful to fix a distinguished path P in T'_n with nodes $r = v_0, v_1, \dots, v_n$ (it may be useful to think of this path as running “along the left-hand side” of T'_n). Each node v_i has one child $v_{i+1} = v_{i+1}^{(0)}$ in P – let its other children be called $v_{i+1}^{(1)}, \dots, v_{i+1}^{(d-1)}$. Denote by $T^{i,j}$ the subtree of T'_n rooted at $v_i^{(j)}$. Let $T_n^{i,j} = T^{i,j} \cap T_n$ (which may be empty), and let $N_{n,m}^{i,j}$ be the set of nodes of $N_{n,m}$ that are contained in $T_n^{i,j}$.

Given an automorphism $\alpha(T'_n)$ of T'_n , we may view α as acting on the labels $X_{v,i}$ and on the permutations σ_v , by viewing the permutations σ_v as fixed to the nodes of T'_n , the labels $X_{v,i}$ as fixed to the edges of T'_n , and both as being moved by the automorphism α . The presence of the permutations σ_v ensures that for any automorphism $\alpha(T'_n)$ of T'_n , the distribution of T'_n (with both permutations and labels attached) is identical to that of $\alpha(T'_n)$ (with both permutations and labels attached), and in particular, α induces a labeled isomorphism between $T_n(T'_n)$ and $T_n(\alpha(T'_n))$. This symmetry, gained by the

addition of the permutations, will greatly simplify later calculations. In particular, it yields the following fact.

Fact 5. *Let E, F , and G be events in the σ -field generated by T'_n with its labeling such that E is invariant under automorphisms of T'_n and there is some automorphism α of T'_n such that $G = \alpha(F)$. Then $\mathbf{P}\{E|F\} = \mathbf{P}\{E|G\}$.*

For example, if F is the event that $S_v \leq m$ for some $m \in \mathbb{R}$ and some given $v \in N_n$, then G could be the event that $S_{v'} \leq m$ for any given $v' \in N_n$. We will often use the phrase “by symmetry” in our arguments, rather than making explicit reference to Fact 5.

1.4 Outline

In Section 2, we prove Theorem 4, and in doing so introduce the idea of *amplification*, which also plays a role in the proof of Theorem 3. In Sections 3 and 4.1, we flesh out the high-level discussion of Reed’s approach given above and explain some of the key ideas behind our proof of Theorem 3, particularly the upper bound on M_n and the importance of leading nodes. In Section 4.2 we prove two key lemmas which allow us to control the shape of random walks conditioned on their value after time n . Finally, in Sections 4.3 and 5, respectively, we prove lower tail bounds and upper tail bounds on M_n that together prove Theorem 3.

2 Proof of Theorem 4

To give an idea of how we will prove Theorem 4, we first consider the special case that the offspring distribution B is deterministically d and that X is deterministically bounded from above, say $X \leq A$ for some constant A . In this case $\mathbf{P}\{S\} = 1$ so the conditioning in Theorem 4 vanishes. We also presume for simplicity that $\text{ess inf } X = 0$. We consider the related branching process T_0 in which the set of children of a node is the set of its children in T for which the displacement is 0. As $\mathbf{P}\{X = 0\} > 1/\mathbf{E}B = 1/d$, T_0 survives with positive probability, so there is a positive probability p_0 that $M_n = 0$ for every n .

Suppose that we want to bound the probability that M_n is greater than x , for some given x . Since $X \leq A$ almost surely, if n is at most x/A then every node at depth n has label at most $nA \leq x$, so $\mathbf{P}\{M_n > x\} = 0$. For larger n , we first observe that for any node v at depth $\lfloor x/A \rfloor$, the tree rooted at v whose nodes are the descendants w of v with $S_w = S_v$ is distributed precisely as T_0 ; we temporarily denote this tree $T_0(v)$.

The tree $T_0(v)$ survives with probability p_0 , and if $T_0(v)$ survives then in particular, there is a descendent w of v at depth n in $T(\mathcal{E})$ for which $S_w = S_v$, so $M_n \leq S_w = S_v \leq x$. It follows that

$$\mathbf{P}\{M_n > x\} \leq \mathbf{P}\left\{ \bigcap_{v \text{ at depth } \lfloor x/A \rfloor} \{T_0(v) \text{ does not survive.}\} \right\}.$$

Since the subtrees rooted at distinct nodes at depth $\lfloor x/A \rfloor$ are independent and there

are $d^{\lfloor x/A \rfloor}$ such nodes, it follows that

$$\begin{aligned} \mathbf{P}\{M_n > x\} &\leq \prod_{v \text{ at depth } \lfloor x/A \rfloor} \mathbf{P}\{T_0(v) \text{ does not survive.}\} \\ &= (1 - p_0)^{d^{\lfloor x/A \rfloor}}, \end{aligned} \quad (12)$$

which in particular proves the tail bound of Theorem 4 and also immediately implies that $\mathbf{E}M_n = O(1)$. The key to the above line of reasoning is the idea of analyzing the subtrees of T rooted at depth $\lfloor x/A \rfloor$ independently in order to strengthen our probability bound. We will hereafter refer to this technique as an *amplification* argument. McDiarmid (1995) uses this idea in much the same fashion as above in his analysis of the minima of branching random walks; it also plays a key role in both (Devroye and Reed, 1995) and (Reed, 2003).

When $\mathbf{P}\{B = d\} < 1$ and X is not necessarily bounded, we can not argue as straightforwardly as above. However, we still have that $\mathbf{P}\{T_0 \text{ survives}\} = p_0$ for some $p_0 > 0$. For $x \geq 0$ integers $d \geq 0$, we temporarily let $F_{x,d}$ be the set of nodes at depth d with label at most x . Using an amplification argument just as we did in deriving (12) immediately yields that for any integers c and n with $c > 0$ and $n \geq d$,

$$\mathbf{P}\{M_n > x \mid |F_x| = c\} \leq (1 - p_0)^c. \quad (13)$$

So, to handle this case, we really need to control the distribution of the number of nodes at a given level of a supercritical branching process whose labels are not too large. To do so, we use the following result of McDiarmid (1995, Lemma 1), who showed that for any supercritical branching process there exist constants $\gamma_0 > 1$, $c_0 > 0$, and $\delta_0 > 0$ such that for all integers $i \geq 1$,

$$\mathbf{P}\{0 < |N_i| \leq \gamma_0^i\} \leq c_0 e^{-\delta_0 i}. \quad (14)$$

Fix γ_0 , c_0 and δ_0 as above, and define the function $\ell(x) = \lceil \log_{\gamma_0} x \rceil$. Since \mathcal{S} occurs precisely if $|N_i| > 0$ for all i , for $n \geq \ell(x)$,

$$\begin{aligned} \mathbf{P}\{M_n > x, \mathcal{S}\} &\leq \mathbf{P}\{0 < |N_{\lfloor \ell(x) \rfloor}| \leq x\} + \mathbf{P}\{M_n > x \mid |N_{\ell(x)}| > x\} \\ &\leq c_0 e^{-\delta_0 x} + \sup_{x < k \leq d^{\ell(x)}} \mathbf{P}\{M_n > x \mid |N_{\ell(x)}| = k\}. \end{aligned} \quad (15)$$

For fixed k in the above range, we have

$$\begin{aligned} \mathbf{P}\{M_n > x \mid |N_{\ell(x)}| = k\} &= \sum_{i=0}^k (\mathbf{P}\{M_n > x \mid |N_{\ell(x)}| = k, |F_{x,\ell(x)}| = i\} \\ &\quad \cdot \mathbf{P}\{|F_{x,\ell(x)}| = i \mid |N_{\ell(x)}| = k\}) \\ &\leq \sum_{i=0}^k (1 - p_0)^i \mathbf{P}\{|F_{x,\ell(x)}| = i \mid |N_{\ell(x)}| = k\} \\ &\leq (1 - p_0)^{k/2} + \mathbf{P}\{|F_{x,\ell(x)}| \leq k/2 \mid |N_{\ell(x)}| = k\}. \end{aligned} \quad (16)$$

Now fix $u > 0$ such that $\mathbf{E}e^{uX} = a < \infty$; such u and a exist by condition (II). We then

have

$$\begin{aligned}
\mathbf{P}\{v_{\ell(x)} \notin F_{x,\ell(x)} \mid v_{\ell(x)} \in N_{\ell(x)}\} &= \mathbf{P}\{x < S_{v_{\ell(x)}} < \infty\} \\
&\leq \frac{[\mathbf{E}e^{uX}]^{\ell(x)}}{e^{ux}} \\
&= e^{(\log a)\lceil \log_{\gamma_0} x \rceil - ux} \\
&\leq c_1 e^{-\delta_1 x},
\end{aligned} \tag{17}$$

for some $c_1 > 0$ and $\delta_1 > 0$ and for all $x > 0$. It follows that

$$\mathbf{E}\{|N_{\ell(x)} \setminus F_{x,\ell(x)}| \mid |N_{\ell(x)}| = k\} \leq c_1 e^{-\delta_1 x} \cdot k,$$

so by Markov's inequality

$$\mathbf{P}\{|F_{x,\ell(x)}| \leq k/2 \mid |N_{\ell(x)}| = k\} \leq 2c_1 e^{-\delta_1 x}. \tag{18}$$

Combining (15), (16) and (18), we thus have

$$\mathbf{P}\{M_n \geq x, \mathcal{S}\} \leq c_2 e^{-\delta_2 x}, \tag{19}$$

for some $c_2 > 0$ and $\delta_2 > 0$ and all $x > 0$. Finally, since T is supercritical, we have $\mathbf{P}\{\mathcal{S}\} = p > 0$. Letting $c_3 = c_2/p$, by (19) we thus have $\mathbf{P}\{M_n > x \mid \mathcal{S}\} \leq c_3 e^{-\delta_2 x}$ for $n \geq \ell(x)$. For $n < \ell(x)$, since given \mathcal{S} there is at least one node in N_n , by symmetry and arguing as in (17) we have

$$\begin{aligned}
\mathbf{P}\{M_n > x \mid \mathcal{S}\} &\leq \mathbf{P}\{x < S_{v_n} < \infty\} \\
&\leq c_1 e^{-\delta_1 x}.
\end{aligned}$$

This proves the exponential tail bounds of Theorem 4 and also shows that $\mathbf{E}\{M_n \mid \mathcal{S}\} = O(1) = \text{ess inf } X + O(1)$. We remark that since we are assuming $\text{ess inf } X = 0$, there is a shorter proof in the case that $n < \ell(x)$ since $M_{\ell(x)} \leq x$ implies $M_n \leq x$. We have given the above argument since it does not use the fact that X is bounded from below, and we will appeal to it when proving Theorem 3.

3 Typical leading nodes

We recall that a node $v \in T_n$ is strictly leading if the random walk ending at v stays strictly above its conditional expected value given S_v , i.e., if it satisfies (9) with strict inequality. For $m \in \mathbb{R}$, we let

$$G_{n,m} = \{v \in N_n : m - 1 \leq S_v \leq m, S_v \text{ is strictly leading}\}. \tag{20}$$

We impose the requirement that $m - 1 \leq S_v$ as it gives us more precise control over the random walk; however, when m is near m^* , Theorem 1 implies that $\mathbf{P}\{S_v \geq m - 1 \mid S_v \leq m\}$ is bounded away from zero. Furthermore, by replacing the constant $m - 1$ by $m - C$ for some large constant C , we could make this probability arbitrarily close to 1. Intuitively, therefore, we can think of $G_{n,m}$ as the set of ‘‘typical’’ leading nodes in N_n with $S_v \leq m$. The following lemma is the promised formalization of (10).

Lemma 6. Given a random walk S with steps distributed as X , and real numbers a, c with $c > \mathbf{E}X \geq a$ and for which $\mathbf{P}\{\mathbf{E}X < X \leq c\} > 0$,

$$\mathbf{P}\{S_n \leq an, S_n \text{ is strictly leading}\} \leq \frac{\mathbf{P}\{S_n \leq an\}}{n}.$$

and

$$\mathbf{P}\{S_n \leq an, S_n \text{ is strictly leading}\} \geq \mathbf{P}\{\mathbf{E}X < X \leq c\} \cdot \frac{\mathbf{P}\{S_{n-1} \leq an - c\}}{n-1}$$

Proof. The proof of Lemma 6 is an adaptation of proofs from Andersen (1953) and Reed (2003). For fixed n , and n' with $n < n' \leq 2n$, let

$$S_{n'} = S_n + S_{n'-n} = \sum_{i=1}^{n'} X_{i(\bmod n)}.$$

We first note that if S_n is strictly leading, then for any $j = 0, \dots, n-1$, the random walk $S^{(j)}$ with $S_i^{(j)} = S_{j+i} - S_j$ for $i = 1, \dots, n$ is not leading, since $S_n^{(j)} = S_n$ and

$$S_{n-j}^{(j)} = S_n - S_j < S_n \left(\frac{n-j}{n} \right).$$

More strongly, an identical argument shows that *at most* one of the random walks $S = S^{(0)}, S^{(1)}, \dots, S^{(n)}$ is strictly leading. Furthermore, as the random variables X_1, \dots, X_n are independent and the event $\{S_n \leq an\}$ is fixed by permutations of X_1, \dots, X_n , it follows that for all $j = 1, \dots, n$,

$$\mathbf{P}\{S_n \leq an, S_n \text{ is strictly leading}\} = \mathbf{P}\left\{S_n \leq an, S_n^{(j)} \text{ is strictly leading}\right\},$$

and so

$$\begin{aligned} n \cdot \mathbf{P}\{S_n \leq an, S_n \text{ is strictly leading}\} &= \mathbf{P}\left\{\bigcup_{j=0}^{n-1} \{S_n \leq an, S_n^{(j)} \text{ is strictly leading}\}\right\} \\ &\leq \mathbf{P}\{S_n \leq an\}, \end{aligned}$$

proving the upper bound of the lemma.

Next, for $i = 1, \dots, n-1$, let $\hat{S}_i = S_{i+1} - X_1$. In order that $S_n \leq an$ and that S_n is *strictly* leading, it suffices that

- $\mathbf{E}X < X_1 \leq c$,
- $\hat{S}_{n-1} \leq an - c$, and
- \hat{S}_{n-1} is leading.

As X_1 is independent from $\hat{S}_1, \dots, \hat{S}_{n-1}$, it follows that

$$\begin{aligned} \mathbf{P}\{S_n \leq an, S_n \text{ is strictly leading}\} &\geq \mathbf{P}\{\mathbf{E}X < X_1 \leq c\} \cdot \\ &\quad \mathbf{P}\left\{\hat{S}_{n-1} \leq an - c, \hat{S}_{n-1} \text{ is leading}\right\} \end{aligned} \quad (21)$$

For $j = 0, \dots, n-2$, define the random walk $\hat{S}^{(j)}$ by $\hat{S}_i^{(j)} = \hat{S}_{j+i} - \hat{S}_j$, for $i = 1, \dots, n-1$ (where, letting $n' = j+i$, if $n' > n-1$ then $\hat{S}_{n'}^{(j)} = \hat{S}_{n-1} + \hat{S}_{n'-(n-1)}$). Again we have that

$\hat{S}_{n-1}^{(j)} = \hat{S}_{n-1}$ for all $j = 0, \dots, n-2$. Furthermore, if \hat{S}_{n-1} is not leading then, letting j^* be an index for which $\hat{S}_{j^*} - \hat{S}_{n-1}(j^*/(n-1))$ is minimized, it follows immediately that for all $i = 1, \dots, n-1$,

$$\begin{aligned} \hat{S}_i^{(j^*)} &= \hat{S}_{i+j^*} - \hat{S}_{j^*} \\ &= \hat{S}_{i+j^*} - \hat{S}_n \left(\frac{i+j^*}{n-1} \right) + \hat{S}_n \left(\frac{i+j^*}{n-1} \right) - \hat{S}_{j^*} \\ &\geq \hat{S}_{j^*} - \hat{S}_n \left(\frac{j^*}{n-1} \right) + \hat{S}_n \left(\frac{i+j^*}{n-1} \right) - \hat{S}_{j^*} \\ &= \hat{S}_n \left(\frac{i}{n-1} \right) = \hat{S}_n^{(j^*)} \left(\frac{i}{n-1} \right), \end{aligned} \quad (22)$$

i.e., $\hat{S}^{(j)}$ is leading at time $n-1$. Therefore, at least one of $\hat{S}^{(0)} = \hat{S}, \hat{S}^{(1)}, \dots, \hat{S}^{(n-2)}$ is leading at time $n-1$. Since X_2, \dots, X_n are independent and the event $\{\hat{S}_{n-1} \leq an-c\}$ is fixed by permutations of X_2, \dots, X_n , we thus have

$$\begin{aligned} \mathbf{P} \left\{ \hat{S}_{n-1} \leq an-c \right\} &= \mathbf{P} \left\{ \hat{S}_{n-1} \leq an-c, \bigcup_{j=0}^{n-2} \{ \hat{S}_{n-1}^{(j)} \text{ is leading} \} \right\} \\ &\leq (n-1) \mathbf{P} \left\{ \hat{S}_{n-1} \leq an-c, \hat{S}_{n-1} \text{ is leading} \right\}. \end{aligned} \quad (23)$$

Combining (21) and (23), it follows that

$$\mathbf{P} \{ S_n \leq an, S_n \text{ is strictly leading} \} \geq \mathbf{P} \{ \mathbf{E}X < X_1 \leq c \} \cdot \frac{\mathbf{P} \left\{ \hat{S}_{n-1} \leq an-c \right\}}{n-1},$$

proving the lower bound of the lemma. \square

By combining the above argument with the bounds from Theorem 1, the following lemma is immediate:

Lemma 7. *Given any function $g(n)$ tending to infinity with n , for any $v \in N_n$ and any m for which $|m^* - m| \leq \sqrt{n}/g(n)$,*

$$\mathbf{P} \{ v \in G_{n,m} \} = \Theta \left(\frac{\mathbf{P} \{ S_v \leq m \}}{n} \right),$$

uniformly over all m in the above range.

We omit the proof of Lemma 7 as it is essentially identical to that of Lemma 6 (taking $a = m/n$ and choosing any fixed $c > \mathbf{E}X$ for which $\mathbf{P} \{ \mathbf{E}X < X \leq c \} > 0$; such c exists since X is non-constant).

4 The upper bound

4.1 A warmup

To prove an upper bound on M_n , we will eventually show that if $\mathbf{E}|G_{n,m}| = \Omega(1)$ then $\mathbf{P} \{|G_{n,m}| > 0\} = \Omega(1)$ (the definition of $G_{n,m}$ appears in (20)). In order to demonstrate one of the key techniques of the lower bound in a simplified setting, we prove:

Lemma 8. For $m > m^*$ for which $m^* - m = o(\sqrt{n})$, if $\mathbf{E}|G_{n,m}| = \Omega(1/n^{5/2})$ then $\mathbf{P}\{|G_{n,m}| \geq 1\} = \Omega(1/n^{5/2}) = \Omega(1/m^{5/2})$.

To prove Lemma 8, we use a version of the second moment method often called the Chung-Erdős inequality (see Chung and Erdős (1952) and also Devroye and Reed (1995) and Alon and Spencer (2000), Chapter 2), which in our setting can be stated as follows: for any integer $i \geq 1$ and any random set $\mathcal{R} \subseteq N'_n$ (recall that N'_n is the set of nodes at depth n in T'_n),

$$\mathbf{P}\{|\mathcal{R}| > 0\} \geq \frac{\mathbf{E}|\mathcal{R}|}{1 + \sup_{v \in N'_n} \mathbf{E}\{|\mathcal{R}| \mid v \in \mathcal{R}\}}. \quad (24)$$

We will apply (24) both here and later in the section. We recall that each node v_i on the distinguished path v_0, \dots, v_{n-1} has children $v_{i+1} = v_{i+1}^{(0)}, v_{i+1}^{(1)}, \dots, v_{i+1}^{(d-1)}$ in T'_n . For $i = 0, \dots, n, j = 0, \dots, d-1$, and $m \in \mathbb{R}$ we hereafter denote

$$N_{n,m}^{i,j} = \{v \in N_{n,m} : v \text{ is a descendent of } v_i^{(j)}\}, \quad (25)$$

and define $G_{n,m}^{i,j}$ similarly.

Proof of Lemma 8. By the symmetry of T'_n , $\mathbf{E}\{|G_{n,m}| \mid v \in G_{n,m}\}$ is identical for all $v \in N'_n$, so letting $\mathcal{R} = G_{n,m}$ in (24), we obtain

$$\mathbf{P}\{|G_{n,m}| \geq 1\} \geq \frac{\mathbf{E}\{|G_{n,m}|\}}{1 + \mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\}}. \quad (26)$$

By symmetry,

$$\begin{aligned} \mathbf{E}\{|G_{n,m}|\} &= d^n \mathbf{P}\{v_n \in G_{n,m}\} \\ &= d^n \mathbf{P}\{v_n \in G_{n,m} \mid v_n \in N_n\} \mathbf{P}\{v_n \in N_n\}. \end{aligned} \quad (27)$$

Given that $v_n \in N_n$, S_{v_n} is just a sum of n i.i.d. random variables distributed as X , so by Lemma 7 and Theorem 1,

$$\begin{aligned} \mathbf{P}\{v_n \in G_{n,m} \mid v_n \in N_n\} &= \Theta\left(\frac{\mathbf{P}\{S_{v_n} \leq m \mid v_n \in N_n\}}{n}\right) \\ &= \Theta\left(\frac{e^{-t(m-m^*)}}{n \cdot [\mathbf{E}B]^d}\right), \end{aligned} \quad (28)$$

where $t < 0$ has been chosen such that $t\Lambda'(t) - \Lambda(t) = \log(\mathbf{E}B)$. Now, for any edge $e = vw$ of the tree T'_n ,

$$\mathbf{P}\{X_e < \infty\} = \sum_{i=0}^d \mathbf{P}\{X_e < \infty \mid B_v = i\} \mathbf{P}\{B_v = i\} = \frac{1}{d} \sum_{i=1}^d i \mathbf{P}\{B_v = i\} = \frac{\mathbf{E}B}{d}. \quad (29)$$

As the variables $\{B_v\}_{v \in T'_n}$ are independent, it follows that for distinct edges $e = vw$ and $f = xy$, X_e and X_f are independent unless $v = x$, i.e., unless w and y are siblings. In particular, it follows from this independence and from (29) that

$$\mathbf{P}\{v_n \in N_n\} = \mathbf{P}\left\{\bigcap_{i=1}^n \{X_{v_{i-1}v_i} < \infty\}\right\} = \left(\frac{\mathbf{E}B}{d}\right)^n. \quad (30)$$

By combining (27), (28), and (30), we obtain the bound

$$\mathbf{E}|G_{n,m}| = \Theta\left(\frac{e^{-t(m-m^*)}}{n}\right). \quad (31)$$

(A brief remark: we will usually omit the sorts of arguments leading to (30) when such calculations arise in later proofs; we have included them once for completeness.) We now turn to bounding $\mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\}$. By symmetry, we have

$$\begin{aligned} \mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\} &\leq \mathbf{E}\{|N_{n,m}| \mid v_n \in G_{n,m}\} \\ &= 1 + \sum_{i=0}^{n-1} \sum_{j=1}^{d-1} \mathbf{E}\{|N_{n,m}^{i+1,j}| \mid v_n \in G_{n,m}\} \\ &= 1 + (d-1) \sum_{i=0}^{n-1} \mathbf{E}\{|N_{n,m}^{i+1,1}| \mid v_n \in G_{n,m}\}. \end{aligned} \quad (32)$$

Since $v_n \in G_{n,m}$, for i with $0 \leq i \leq n$ we have $S_{v_i} \geq (m-1)(i/n) \geq mi/n - 1$. It follows that for such i , $N_{n,m}^{i+1,1}$ is at most the number of descendants v of $v_{i+1}^{(1)}$ for which $S_v - S_{v_i}$ is at most $m(n-i)/n + 1$. We recall that $m_i^* = \Lambda'(t)i - \log i/2t$. Let $\Delta_i = mi/n + 1 - m_i^*$ for $i = 1, \dots, n$, and fix some distinguished descendant v of $v_{i+1}^{(1)}$ at depth n . By the previous observation, linearity of expectation and symmetry we have

$$\begin{aligned} \mathbf{E}\{|N_{n,m}^{i+1,1}| \mid v_n \in G_{n,m}\} &\leq |N_n^{i+1,1}| \mathbf{P}\{S_v - S_{v_i} \leq m_{n-i}^* + \Delta_{(n-i)} \mid v_n \in G_{n,m}\} \\ &= d^{n-(i+1)} \cdot \left(\frac{\mathbf{E}B}{d}\right)^{n-(i+1)} \cdot \\ &\quad \mathbf{P}\{S_v - S_{v_i} \leq m_{n-i}^* + \Delta_{(n-i)} \mid v \in N_n^{i+1,1}\} \\ &= (\mathbf{E}B)^{n-(i+1)} \cdot \Theta\left(\frac{e^{-t\Delta_{n-i}}}{(\mathbf{E}B)^{n-i}}\right) = \Theta(e^{-t\Delta_{n-i}}), \end{aligned} \quad (33)$$

We now claim that for all $i = 1, \dots, n$,

$$e^{-t\Delta_{n-i}} = O(n^{1/2}e^{-t(m-m^*)}). \quad (34)$$

From (32), (33), and (34), it follows that

$$\mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\} \leq 1 + O(n^{3/2}e^{-t(m-m^*)}). \quad (35)$$

Combining (31) and (35) we see that if $\mathbf{E}|G_{n,m}| = \Omega(1/n^{5/2})$ then $\mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\} \leq 1 + O(n^{5/2} \cdot \mathbf{E}|G_{n,m}|)$, which together with (27) proves the fact. To see that (34) holds,

we write

$$\begin{aligned}
\Delta_{n-i} &= \frac{m(n-i)}{n} - m_{n-i}^* + 1 \\
&= \frac{m(n-i)}{n} - \Lambda'(t)(n-i) - \frac{\log(n-i)}{2t} + 1 \\
&= m - \frac{mi}{n} + \Lambda'(t)i - \Lambda'(t)n - \frac{\log n}{2t} + \frac{\log n - \log(n-i)}{2t} + 1 \\
&= m - m^* + \Lambda'(t)i - \frac{mi}{n} + \frac{\log n - \log(n-i)}{2t} + 1 \\
&< m - m^* + \frac{i}{n} (\Lambda'(t)n - m) + 1 \\
&\leq m - m^* - \frac{\log n}{2t} + 1,
\end{aligned}$$

the final inequality holding as $m \geq m^* = \Lambda'(t)n + \log n/(2t)$ (recall that t is negative). Thus $\Delta_{n-i} \leq m - m^* - \log n/2t$, from which (34) follows immediately. \square

By Lemma 8 and a standard amplification argument, we could show that $\mathbf{P}\{|N_{n,m+a \log n}| \geq 1\} = \Omega(1)$ for some $a = O(1)$ and thereby show that $\mathbf{E}\{M_n \mid \mathcal{S}\} \leq m^* + a \log n$; we will not do so as we are headed towards stronger upper bounds, the key step of which is to strengthen Lemma 8. We now state this aim formally.

Lemma 9. *For all values of m for which $m^* - m = o(\sqrt{n})$,*

$$\mathbf{E}\{|G_{n,m}| \mid v_n \in G_{n,m}\} = \mathbf{E}|G_{n,m}| + O(1). \quad (36)$$

Assume for the moment that Lemma 9 holds, and let

$$m' = m^* - \frac{\log n}{2t} = \Lambda'(t)n - \frac{3 \log n}{2t}.$$

By Lemma 7 and linearity of expectation, we see that m' is within $O(1)$ of the smallest value m for which $\mathbf{E}|G_{n,m}| \geq 1$. Applying (24) with $\mathcal{R} = G_{n,m'}$, it follows immediately that $\mathbf{P}\{|G_{n,m'}| > 0\} = \Omega(1)$. Since $G_{n,m} \subseteq N_{n,m}$ for all m , it follows that $\mathbf{P}\{|N_{n,m'}| > 0\} = \Omega(1)$, so $\mathbf{P}\{M_n \leq m'\} = \Omega(1)$. By applying an amplification argument exactly as we did when proving Theorem 4, we immediately obtain exponential tail bounds for the upper tail of M_n :

Corollary 10. *There exist constants $C_1 > 0, \delta_1 > 0$ such that for all $x > 0$,*

$$\mathbf{P}\{M_n \geq m' + x \mid \mathcal{S}\} \leq C_1 e^{-\delta_1 x}.$$

To obtain the upper tail bounds of Theorem 3, it thus remains to prove Lemma 9.

4.2 The shape of random walks

The proof of Lemma 9 is based on establishing finer control over the behavior of a given random walk in T than that given by Lemma 7. In this section, we prove the two lemmas that accomplish this. In Lemma 11, we bound the probability that a walk S_0, S_1, \dots, S_n is ever very far below its conditional mean given S_n ; in Lemma 12, we bound the probability that such a random walk is ever *close to* its conditional

mean, given that it is *never* very far below its conditional mean. These results are rather straightforward; the only technicalities are due to the fact that we must apply Theorem 1 in the course of the proofs. We now proceed to the details.

For $a \leq 0$, we say that S *stays above* a up to time n , and write $\{S_n \text{ abo } a\}$, if

$$S_i > S_n \cdot \frac{i}{n} + a, \text{ for all } i = 1, \dots, n.$$

In particular, $\{S_n \text{ abo } 0\}$ is simply the event that S_n is strictly leading.

Lemma 11. *Given any function $g(n)$ tending to plus infinity with n , any m for which $|\Lambda'(t)n - m| = \sqrt{n}/g(n)$, and any $a \leq -1$,*

$$\mathbf{P} \{S_n \text{ abo } a \mid m - 1 \leq S_n \leq m\} = O\left(\frac{|a|^9}{n}\right), \quad (37)$$

uniformly over all m and a in the above ranges.

Proof. For simplicity we assume that a is an integer, that $m = \Lambda'(t)n$, and that $g(n) \leq \log n$; this eases the notational burden without changing the essence of the proof. The probability in (37) increases as a decreases; it thus suffices to prove (37) when $|a|$ is at least some large fixed constant C . We assume $|a|$ is large enough that $|a| \leq \sqrt{|a|^3/\log(|a|^3)} \leq \sqrt{|a|}/g(|a|)$.

We remark that if $|a| > n^{1/9}$ then (37) holds trivially; we thus assume that $|a| \leq n^{1/9}$. Let $n_1 = n + 2|a|^3$, and let $m_1 = \Lambda'(t)n_1$; as $|a| \leq n^{1/9}$, we have $n_1 = \Theta(n)$. Let S' be the random walk with $S'_i = S_{|a|^3+i} - S_{|a|^3}$ (the original walk “started at time $|a|^3$ ”). The walk S' is distributed as S ; we will show that

$$\mathbf{P} \{S'_n \text{ abo } a \mid m - 1 \leq S'_n \leq m\} = O\left(\frac{|a|^9}{n}\right), \quad (38)$$

which proves the lemma. We proceed by comparing the following two probabilities:

$$\mathbf{P} \{m_1 - 3 \leq S_{n_1} \leq m_1, S_{n_1} \text{ abo } 0\} \quad \text{and} \quad \mathbf{P} \{m - 1 \leq S'_n \leq m, S'_n \text{ abo } a\}.$$

By Lemma 7,

$$\begin{aligned} \mathbf{P} \{m_1 - 3 \leq S_{n_1} \leq m_1, S_{n_1} \text{ abo } 0\} &= \Theta\left(\frac{\mathbf{P} \{m_1 - 3 \leq S_{n_1} \leq m_1\}}{n_1}\right) \\ &= \Theta\left(\frac{\mathbf{P} \{m_1 - 3 \leq S_{n_1} \leq m_1\}}{n}\right) \end{aligned} \quad (39)$$

By Theorem 1,

$$\begin{aligned} \mathbf{P} \{m_1 - 3 \leq S_{n_1} \leq m_1\} &= \Theta\left(\frac{e^{-\Lambda'(t)m_1}}{\sqrt{n_1}}\right) = \Theta\left(\frac{e^{-2|a|^3\Lambda'(t)}e^{-\Lambda'(t)n}}{\sqrt{n}}\right) \\ &= \Theta\left(e^{-2|a|^3\Lambda'(t)}\mathbf{P} \{m - 1 \leq S'_n \leq m\}\right), \end{aligned} \quad (40)$$

so, letting E^* be the event $\{m_1 - 3 \leq S_{n_1} \leq m_1, S_{n_1} \text{ abo } 0\}$, (39) and (40) yield that

$$\mathbf{P} \{E^*\} = \Theta\left(\frac{e^{-2|a|^3\Lambda'(t)}\mathbf{P} \{m - 1 \leq S'_n \leq m\}}{n}\right). \quad (41)$$

We will show that

$$\mathbf{P}\{E^*\} = \Omega\left(\frac{e^{-2|a|^3\Lambda'(t)}\mathbf{P}\{m-1 \leq S'_n \leq m, S'_n \text{ abo } a\}}{|a|^9}\right). \quad (42)$$

Equation (38) follows immediately from (41) and (42); it thus remains to prove (42).

We claim that for E^* to occur it suffices that the following events occur:

- (1) $\Lambda'(t)|a|^3 + |a| + 1 \leq S_{|a|^3} \leq \Lambda'(t)|a|^3 + |a| + 2$, which we denote E_1 ;
- (2) $\Lambda'(t)|a|^3 - (|a| + 3) \leq S_{n_1} - S_{n_1-|a|^3} \leq \Lambda'(t)|a|^3 - (|a| + 2)$, which we denote E_2 ;
- (3) $m - 1 \leq S'_n \leq m$, which we denote E_3 ;
- (4) $S_{|a|^3} \text{ abo } 0$;
- (5) letting $S_j^{(1)} = S_{n_1-|a|^3+j} - S_{n_1-|a|^3}$ for $j = 0, 1, \dots, |a|^3$, $S_{|a|^3}^{(1)} \text{ abo } 0$; and finally,
- (6) $S'_n \text{ abo } a$.

Informally, (1) and (4) ensure that the first $|a|^3$ steps of the walk do not prevent E^* from occurring, (2) and (5) do the same for the last $|a|^3$ steps of the walk, and (3) and (6) do likewise for the middle $n_1 - 2|a|^3 = n$ steps of the walk. (This is not quite the whole story; the “extra height” (between $|a| + 1$ and $|a| + 2$) gained in (1), as well as the extra height gained in (2), are required so that (3) and (6) can do their job.)




Figure 1: The events (1)-(6). For example, (1) occurs because after $|a|^3$ steps the walk is between the lowest two dashed horizontal lines, and (4) occurs because in the first $|a|^3$ steps the walk stays above the dashed line segment connecting its endpoints. The event E^* occurs because after n_1 steps the walk is between $m_1 - 3$ and m_1 and has stayed above dashed line connecting its endpoints.

By the independence of disjoint sections of the random walk, we thus have

$$\begin{aligned} \mathbf{P}\{E^*\} &\geq \mathbf{P}\{E_1, E_2, E_3, S_{|a|^3} \text{ abo } 0, S_{|a|^3}^{(1)} \text{ abo } 0, S'_n \text{ abo } a\} \\ &= \mathbf{P}\{E_1, S_{|a|^3} \text{ abo } 0\} \cdot \mathbf{P}\{E_2, S_{|a|^3}^{(1)} \text{ abo } 0\} \cdot \mathbf{P}\{E_3, S'_n \text{ abo } a\}. \end{aligned} \quad (43)$$

As $|a| \leq \sqrt{|a|^3}/g(|a|)$, we may apply Theorem 1 and Lemma 7 to bound $\mathbf{P}\{E_1, S_{|a|^3} \text{ abo } 0\}$:

$$\mathbf{P}\{E_1, S_{|a|^3} \text{ abo } 0\} = \Theta\left(\frac{\mathbf{P}\{E_1\}}{|a|^3}\right) = \Theta\left(\frac{e^{-|a|^3\Lambda'(t)}e^{-t|a|}}{|a|^3\sqrt{|a|^3}}\right) = \Theta\left(\frac{e^{-|a|^3\Lambda'(t)}e^{-t|a|}}{|a|^{9/2}}\right). \quad (44)$$

Likewise, we have

$$\mathbf{P}\{E_2, S_{|a|^3}^{(1)} \text{ abo } 0\} = \Theta\left(\frac{\mathbf{P}\{E_2\}}{|a|^3}\right) = \Theta\left(\frac{e^{-|a|^3\Lambda'(t)}e^{t|a|}}{|a|^{9/2}}\right), \quad (45)$$

Combining (43), (44), and (45) yields that

$$\mathbf{P}\{E^*\} = \Omega\left(\frac{e^{-2|a|^3\Lambda'(t)}}{|a|^9}\mathbf{P}\{E_3, S'_n \text{ abo } a\}\right),$$

which is precisely the claim in (42). \square

Using Lemma 11, we can bound the conditional probability that S spends much time *near* its mean, given that it is never much below its conditional mean up to time n . More precisely: for $0 \leq k \leq n$, let $b(n, k) = \min\{k, n - k\}$, and let C_k be the event that $S_k \leq S_n(k/n) + b(n, k)^{1/57}$ (that S_k is “close” to its conditional mean; the meaning of “close” depends on how near k is to one of the ends of the random walk). Given an integer $a \leq -1$, let $B_a = \bigcup_{k=|a|^{57}}^{n-|a|^{57}} C_k$ (if $|a|^{57} > n/2$ then B_a is the empty event). Then:

Lemma 12. *Given any function $g(n)$ tending to plus infinity with n , for any m for which $|\Lambda'(t)n - m| \leq \sqrt{n}/g(n)$ and any integer $a \leq -1$,*

$$\mathbf{P}\{S_n \text{ abo } a, B_a \mid m - 1 \leq S_n \leq m\} = O\left(\frac{1}{n|a|^9}\right). \quad (46)$$

Proof. As in the proof of Lemma 11, for the sake of readability we assume that $m = \Lambda'(t)n$ and that $g(k) \leq \log k$ for all k . We may presume that $|a|^{57} < n/2$ or else the claim holds trivially. For any fixed K , for all a with $|a| \leq K$, the claim follows immediately from Lemma 11, so we may assume $|a|$ is larger than any fixed K ; we henceforth assume $|a|$ is large enough that $|a| \leq \sqrt{|a|^{57}/g(|a|)}$.

Choose k with $b(n, k) \geq |a|^{57}$. If $\{m - 1 \leq S_n \leq m\}$ and C_k are to both occur then necessarily

$$S_k \leq \frac{mk}{n} + b(n, k)^{1/57} = \Lambda'(t)k + b(n, k)^{1/57}. \quad (47)$$

Similarly, if $\{m - 1 \leq S_n \leq m\}$ and $\{S_n \text{ abo } a\}$ are to both occur then necessarily

$$S_k \geq \Lambda'(t)k + a - 1 \quad (48)$$

(recall that a is negative). More strongly, for C_k , $\{m - 1 \leq S_n \leq m\}$ and $\{S_n \text{ abo } a\}$ to *all* occur, there must be some integer i with $a - 1 \leq i \leq b(n, k)^{1/57}$ for which the following four events occur:

- $\Lambda'(t)k + i \leq S_k \leq \Lambda'(t)k + i + 1$, which we denote $A_{i,k}$;
- $\{S_k \text{ abo } a - (i + 1)\}$, which we denote $D_{i,k}$;

and letting $S'_j = S_{k+j} - S_j$ for $0 \leq j \leq n - k$,

- $\Lambda'(t)(n - k) - (i + 1) \leq S'_{n-k} \leq \Lambda'(t)(n - k) - i$, which we denote $E_{i,k}$;
- $\{S'_{n-k} \text{ abo } a - (i + 1)\}$, which we denote $F_{i,k}$.

Restating the above using the names of these events, we have

$$C_k \cap \{S_n \text{ abo } a\} \cap \{m - 1 \leq S_n \leq m\} \subset \bigcup_{i=a-1}^{\lfloor b(n,k)^{1/57} \rfloor} A_{i,k} \cap D_{i,k} \cap E_{i,k} \cap F_{i,k}. \quad (49)$$

Since $B_a = \bigcup_{k=|a|}^{n-|a|} C_k$, It follows by a union bound that

$$\mathbf{P} \{B_a, S_n \text{ abo } a, m - 1 \leq S_n \leq n\} \leq \sum_{k=|a|}^{n-|a|} \sum_{i=a-1}^{\lfloor b(n,k)^{1/57} \rfloor} \mathbf{P} \{A_{i,k}, D_{i,k}, E_{i,k}, F_{i,k}\}. \quad (50)$$

We will show that for each k with $b(n, k) \geq |a|^{57}$, for all i in the above range,

$$\mathbf{P} \{A_{i,k}, D_{i,k}, E_{i,k}, F_{i,k}\} = O \left(\mathbf{P} \{m - 1 \leq S_n \leq m\} \cdot \frac{b(n, k)^{18/57} n^{1/2}}{(k(n - k))^{3/2}} \right). \quad (51)$$

We remark that the bound in (51) does not depend on i . Presuming for a moment that (51) holds, by dividing through by $\mathbf{P} \{m - 1 \leq S_n \leq m\}$ in (50) and (51), we obtain

$$\begin{aligned} \mathbf{P} \{B_a, S_n \text{ abo } a \mid m - 1 \leq S_n \leq n\} &= \sum_{k=|a|}^{n-|a|} \sum_{i=a-1}^{\lfloor b(n,k)^{1/57} \rfloor} \frac{\mathbf{P} \{A_{i,k}, D_{i,k}, E_{i,k}, F_{i,k}\}}{\mathbf{P} \{m - 1 \leq S_n \leq m\}} \\ &= O \left(\sum_{k=|a|}^{n-|a|} \sum_{i=a-1}^{\lfloor b(n,k)^{1/57} \rfloor} \frac{b(n, k)^{18/57} n^{1/2}}{(k(n - k))^{3/2}} \right) \end{aligned} \quad (52)$$

Since $b(n, k)^{1/57} \geq |a|$, the inner sum in (52) has at most $2b(n, k)^{1/57} + 2$ identical terms. We thus have

$$\begin{aligned} \mathbf{P} \{B_a, S_n \text{ abo } a \mid m - 1 \leq S_n \leq n\} &= O \left(\sum_{k=|a|}^{n-|a|} \frac{b(n, k)^{19/57} n^{1/2}}{(k(n - k))^{3/2}} \right) \\ &= O \left(\sum_{k=|a|}^{\lceil n/2 \rceil} \frac{b(n, k)^{19/57} n^{1/2}}{(k(n - k))^{3/2}} \right) \\ &= O \left(\frac{1}{n} \sum_{k=|a|}^{\lceil n/2 \rceil} \frac{k^{19/57}}{k^{3/2}} \right) \\ &= O \left(\frac{1}{n} \sum_{k=|a|}^{\lceil n/2 \rceil} \frac{1}{k^{7/6}} \right) = O \left(\frac{1}{|a|^9 \cdot n} \right), \end{aligned}$$

which proves the lemma. It therefore remains to prove (51). To do so, we first write

$$\mathbf{P} \{A_{i,k}, D_{i,k}\} = \mathbf{P} \{A_{i,k}\} \cdot \mathbf{P} \{D_{i,k} \mid A_{i,k}\}.$$

Since $|i| \leq \lfloor b(n, k) \rfloor^{1/57} \leq k^{1/57} \leq \sqrt{k}/g(k)$, and likewise $|a| \leq b(n, k)^{1/57}$, we have $|a - (i + 1)| \leq 2\sqrt{k}/g(k) + 1$. By Lemma 11 (applied with a slightly different function g) it then follows that

$$\mathbf{P}\{D_{i,k} \mid A_{i,k}\} = O\left(\frac{(|a| + |i| + 1)^9}{k}\right) = O\left(\frac{b(n, k)^{9/57}}{k}\right), \quad (53)$$

so

$$\mathbf{P}\{A_{i,k}, D_{i,k}\} = O\left(\frac{\mathbf{P}\{A_{i,k}\} \cdot b(n, k)^{9/57}}{k}\right), \quad (54)$$

and an identical derivation yields

$$\mathbf{P}\{E_{i,k}, F_{i,k}\} = O\left(\frac{\mathbf{P}\{E_{i,k}\} \cdot b(n, k)^{9/57}}{n - k}\right). \quad (55)$$

By the independence of disjoint sections of the random walk, $A_{i,k}$ and $B_{i,k}$ are independent of $E_{i,k}$ and $F_{i,k}$, so $\mathbf{P}\{A_{i,k}, D_{i,k}, E_{i,k}, F_{i,k}\} = \mathbf{P}\{A_{i,k}, D_{i,k}\} \mathbf{P}\{E_{i,k}, F_{i,k}\}$, and by (54) and (55) we thus have

$$\mathbf{P}\{A_{i,k}, D_{i,k}, E_{i,k}, F_{i,k}\} = O\left(\frac{\mathbf{P}\{A_{i,k}\} \mathbf{P}\{E_{i,k}\} \cdot b(n, k)^{18/57}}{k(n - k)}\right). \quad (56)$$

Finally, the events $A_{i,k}$, $E_{i,k}$, and $\{m - 1 \leq S_n \leq m\}$ all simply restrict the value of a certain random walk at its endpoint; by applying Theorem 1 to bound $\mathbf{P}\{A_{i,k}\}$, $\mathbf{P}\{E_{i,k}\}$, and $\mathbf{P}\{m - 1 \leq S_n \leq m\}$, it is easily seen that

$$\mathbf{P}\{A_{i,k}\} \cdot \mathbf{P}\{E_{i,k}\} = \Theta\left(\mathbf{P}\{m - 1 \leq S_n \leq m\} \cdot \frac{n^{1/2}}{(k(n - k))^{1/2}}\right).$$

Combining this last equation with (56) proves (51) and completes the proof. \square

It is interesting to compare Lemmas 11 and 12 with the work of Bramson (1978b) in studying branching Brownian motion; he required quite similar bounds on the behavior of a Brownian bridge (or equivalently, of Brownian motion conditioned on its value after some time t). It would be interesting to find a proof of Theorem 3, or at least of Lemmas 11 and 12, that proceeded via comparison with Brownian motion. However, such an approach may be quite difficult, given the lack of success to date at transferring Bramson's argument to a discrete setting. At any rate, with these lemmas under our belt we are ready to prove Lemma 9.

4.3 Proof of Lemma 9

The chain of reasoning is similar to that in the proof of Lemma 8. We first recall that for a node v at depth n , by Lemma 7

$$\mathbf{P}\{v \in G_{n,m'}\} = \Theta\left(\frac{\mathbf{P}\{m' - 1 \leq S_v \leq m'\}}{n}\right). \quad (57)$$

Recalling the event B_a defined just before Lemma 12, we will write $B_a(v)$ for the event that B_a occurs for the random walk ending at v . Since $|\Lambda'(t)n - m'| = O(\log n)$, by Lemmas 11 and 12, for all integers $a < -1$,

$$\begin{aligned} \mathbf{P} \{m' - 1 \leq S_v \leq m, S_v \text{ abo } a, B_a(v)\} &= \Theta \left(\frac{\mathbf{P} \{m' - 1 \leq S_v \leq m\}}{n|a|^9} \right) \\ &= O \left(\frac{\mathbf{P} \{v \in G_{n,m'}\}}{|a|^9} \right). \end{aligned}$$

It follows that there is some large constant $C > 0$ such that if $v \in G_{n,m'}$ then, letting R_0, R_1, \dots, R_n be the partial sums on the path to v , with probability at least $1/2$

$$R_k \geq R_n \cdot \frac{k}{n} + b(n, k)^{1/57} \text{ for all } C \leq k \leq n - C. \quad (58)$$

If $v \in G_{n,m'}$ and v additionally satisfies (58), we say that v is *well-behaved*. We write $W_{n,m'}$ for the set of well-behaved nodes in $G_{n,m'}$. We emphasize that $W_{n,m'} \subseteq G_{n,m'} \subseteq N_{n,m'}$. Furthermore, since each node in $G_{n,m'}$ is well-behaved with probability at least $1/2$, it follows from the definition of m' that $\mathbf{E}|W_{n,m'}| = \Omega(\mathbf{E}|G_{n,m'}|) = \Omega(1)$. We claim that

$$\mathbf{E} \{|W_{n,m'}| \mid v_n \in W_{n,m'}\} = \mathbf{E}|W_{n,m'}| + O(1). \quad (59)$$

Applying (24) with $\mathcal{R} = W_{n,m'}$, it immediately follows that $\mathbf{P} \{|W_{n,m'}| \geq 1\} = \Omega(1)$, so since $W_{n,m'} \subseteq N_{n,m'}$ we have $\mathbf{P} \{|N_{n,m'}| \geq 1\} = \Omega(1)$ as claimed. We thus turn to proving (59).

We now remind the reader of some notation from earlier in the section, and introduce a few new terms. We recall that node v_{i-1} has child $v_i^{(0)} = v_i$ that is on the distinguished path P , and that its remaining children are $v_i^{(1)}, \dots, v_i^{(d-1)}$. Node $v_i^{(j)}$ is the root of a subtree of T_n that we shall call $T_n^{i,j}$. We let $W_{n,m}^{i,j}$ be subset of $W_{n,m}^{i,j}$ contained in $T_n^{i,j}$; this is consistent with the notation $G_{n,m}^{i,j}$ and $N_{n,m}^{i,j}$ introduced in and just after (25). For each $0 \leq i \leq n-1$ fix an arbitrary node x_i at depth n that is a descendent of $v_{i+1}^{(1)}$, and let the partial sums of the labels on the path from v_i to x_i be $S_{x_0,0} = 0, S_{x_i,1}, S_{x_i,2}, \dots, S_{x_i,n-i}$. We remark that the edge labels contributing to the sum $S_{x_i,n-i}$ are a subset of the edge labels contributing to the vertex label S_{x_i} ; more precisely,

$$S_{x_i} = S_{x_i,n-i} + S_{v_i}. \quad (60)$$

Finally, to simplify notation, for $0 \leq i \leq n-1$ let $W_i = W_{n,m'}^{i,1}$, and denote the functions $\mathbf{P} \{\cdot \mid v_n \in W_{n,m'}\}$ and $\mathbf{E} \{\cdot \mid v_n \in W_{n,m'}\}$ by $\mathbf{P}^w \{\cdot\}$ and $\mathbf{E}^w \{\cdot\}$, respectively. We now mimic the portion of the proof of Lemma 8 that leads to (32), in our case for the particular value $m = m'$. By symmetry, we have

$$\begin{aligned} \mathbf{E}^w \{|W_{n,m'}|\} &= 1 + \sum_{i=1}^n \sum_{j=1}^{d-1} \mathbf{E}^w \{W_{n,m'}^{i,j}\} \\ &= 1 + \sum_{i=1}^n (d-1) \mathbf{E}^w \{W_i\}. \end{aligned} \quad (61)$$

Since d does not depend on n and, for a given i , $W_{h'-i} \leq i^d$, it follows that for any fixed integer $0 < c = O(1)$, $\sum_{i=n-c}^n (d-1) \mathbf{E}^w \{W_i\} \leq \sum_{i=1}^c (d-1) i^d = O(1)$. By this

fact, by (61), and since $(d-1)$ is constant, to prove the lemma it therefore suffices to show that

$$\sum_{i=1}^{h'-C} \mathbf{E}^w \{|W_i|\} = O(1), \quad (62)$$

where C is the same constant as in (58).

By symmetry, $\mathbf{E}^w \{|W_i|\} = d^{n-i} \mathbf{P}^w \{x_i \in W_i\}$. In order for $x_i \in W_i$ to occur, we must in particular have that $S_{x_i} \leq m'$, so by (60), we must have $S_{x_i, n-i} \leq m' - S_{v_i}$. The fact that v_n is well-behaved allows us to bound $m' - S_{v_i}$ from above. Furthermore, $S_{x_i, n-i}$ is distributed as $S_{v_{n-i}}$ and is independent of S_{v_i} . This independence will allow us use Theorem 1 to bound the conditional probability that $S_{x_i, n-i} \leq m' - S_{v_i}$. When i is far from 1 and from $n-C$, the bounds on S_i given by the fact that v_n is well-behaved will ensure that S_{v_i} is large enough that the conditional probability that $S_{x_i, n-i} \leq m' - S_{v_i}$ is extremely small. By slightly modifying this same approach, we will prove similar bounds when i is near 1 or near $n-C$; summing these bounds will prove (62). We now turn to the details.

Fix some function g with $g(k) \leq \log k$ for all k and with $g(k)$ tending to plus infinity as k tends to infinity. For any i , for all c with $|c| \leq \sqrt{n-i}/g(i)$, by Theorem 1 and by the independence of X_{v_1}, \dots, X_{v_n} from $S_{x_i, n-i}$, we have

$$\begin{aligned} \mathbf{P}^w \{S_{x_i, n-i} \leq \Lambda'(t)(n-i) + c\} &= \mathbf{P} \{S_{x_i, n-i} \leq \Lambda'(t)(n-i) + c \mid S_{x_i, n-i} < \infty\} \\ &\quad \cdot \mathbf{P} \{S_{x_i, n-i} < \infty\} \\ &= \Theta \left(\frac{e^{-tc}}{\sqrt{n}} \cdot \frac{1}{d^{n-i}} \right) \end{aligned} \quad (63)$$

Furthermore, by (60),

$$\mathbf{P}^w \{S_{x_i} \leq m'\} = \mathbf{P}^w \{S_{x_i, n-i} \leq m' - S_i\}. \quad (64)$$

Let $r = r(n) = \lceil (2 \log n / |t|)^{57} \rceil$.

Case 1 ($r \leq i \leq n-r$): for any $k \geq C$, since $v_n \in W_{n, m'}$, we have $S_{v_k} \geq m' \cdot (k/n) - 1 + b(n, k)^{1/57}$, so

$$m' - S_{v_i} \leq m' \cdot \left(\frac{n-i}{n} \right) + 1 - b(n, i)^{1/57} \quad (65)$$

$$\leq \Lambda'(t)(n-i) + \frac{3 \log n}{2|t|} - b(n, i)^{1/57}. \quad (66)$$

(We will also use (65) in the case that $i < n-r$). When $b(n, i)^{1/57} \geq 2 \log n / |t|$, certainly $\log n / 2|t| \leq \sqrt{n-i}/g(n-i)$, so by (63), (64), and (66) we have

$$\mathbf{P}^w \{S_{x_i} \leq m'\} \leq \mathbf{P}^w \left\{ S_{x_i, n-i} \leq \Lambda'(t)(n-i) - \frac{\log n}{2|t|} + 1 \right\} = \Theta \left(\frac{1}{d^{n-i} n} \right), \quad (67)$$

so by linearity of expectation and symmetry, $\mathbf{E}^w |W_i| = O(1/n)$ for such i . Letting $r = r(n) = \lceil (2 \log n / |t|)^{57} \rceil$, then, we have

$$\sum_{i=r}^{n-r} \mathbf{E}^w |W_i| = O \left(\frac{n-2r}{n} \right) = O(1).$$

This bounds the bulk of the sum (62); it remains to consider the cases when i is either close to 1 or close to $n - C$.

Case 2 ($n - r \leq i \leq n - C$): let $k = n - i$, so $C \leq k < r$. Since $k = O((\log n)^{57})$, $m' \cdot (k/n) = \Lambda'(t)k + O(1)$, so from (64), (65), and the fact that $b(n, i) = k$ we have

$$\begin{aligned} \mathbf{P}^w \{x_i \in W_{n,m'}\} &\leq \mathbf{P}^w \{S_{x_i} \leq m'\} = \mathbf{P} \{S_{x_i, n-i} \leq m' - S_{v_i}\} \\ &= O(\mathbf{P} \{S_{x_i, n-i} \leq \Lambda'(t)k - b(n, i)^{1/57}\}) \\ &= O(\mathbf{P} \{S_{x_i, n-i} \leq \Lambda'(t)k - k^{1/57}\}), \end{aligned}$$

which together with (63) gives

$$\mathbf{P}^w \{x_i \in W_{n,m'}\} = O\left(\frac{e^{tk^{1/57}}}{d^k}\right).$$

By linearity of expectation and by symmetry, it follows that $\mathbf{E}^w |W_i| = O(e^{tk^{1/57}}) = O(k^{-3})$ (recall that t is negative). Therefore

$$\sum_{i=n-r}^{n-C} \mathbf{E}^w |W_i| = O\left(\sum_{k=C}^r \frac{1}{k^3}\right) = O(1).$$

Case 3 ($1 \leq i \leq r$): Let $\Delta_i = S_{v_i} - m' \cdot (i/n)$. Since $v_n \in G_{n,m'}$, necessarily $\Delta_i \geq -i/n \geq -1$. In order that $x_i \in G_{n,m'}$, it is necessary that

- $m' \cdot ((n - i)/n) - (\Delta_i + 1) \leq S_{x_i, n-i} \leq m' \cdot ((n - i)/n) - \Delta_i$ (call this event E_i), and that
- $S_{x_i, n-i} \text{ abo } - (\Delta_i + 1)$,

so

$$\mathbf{P}^w \{x_i \in W_{n,m'}\} \leq \mathbf{P}^w \{x_i \in G_{n,m'}\} \leq \mathbf{P}^w \{E_i, S_{x_i, n-i} \text{ abo } - (\Delta_i + 1)\}.$$

We shall show that

$$\mathbf{P}^w \{E_i, S_{x_i, n-i} \text{ abo } - (\Delta_i + 1)\} = O\left(\frac{1}{d^{n-i}(n-i)}\right), \quad (68)$$

from which it follows just as above that $\sum_{i=1}^r \mathbf{E}^w |W_i| = O(1)$. It thus remains to prove (68). We first observe that by the same argument used to prove (67), we have

$$\begin{aligned} \mathbf{P}^w \left\{ E_i, S_{x_i, n-i} \text{ abo } - (\Delta_i + 1) \mid \Delta_i \geq \frac{2 \log n}{|t|} \right\} &\leq \mathbf{P}^w \left\{ E_i \mid \Delta_i \geq \frac{2 \log n}{|t|} \right\} \\ &\leq \mathbf{P} \left\{ S_{x_i, n-i} \leq \Lambda'(t)(n-i) - \frac{\log n}{2|t|} \right\} \\ &= O\left(\frac{1}{d^{n-i} \cdot n}\right). \end{aligned} \quad (69)$$

Furthermore, if $\Delta_i < 2 \log n / |t|$, then $\Delta_i \leq \sqrt{n-i}/g(n-i)$, so by applying first Lemma 11, then Theorem 1, we obtain

$$\begin{aligned} \mathbf{P}^w \left\{ E_i, S_{x_i, n-i} \text{ abo } - (\Delta_i + 1) \mid \Delta_i < \frac{2 \log n}{|t|} \right\} &= O\left(\frac{\Delta_i^9 \mathbf{P}^w \{E_i \mid \Delta_i < (2 \log n)/|t|\}}{n-i}\right) \\ &= O\left(\frac{\Delta_i^9 e^{\Delta_i t}}{n-i}\right) = O\left(\frac{1}{d^{n-i}(n-i)}\right) \end{aligned} \quad (70)$$

Combining (69) and (70) proves (68) and completes the proof of Lemma 9.

5 The lower bound

We know that the naive approach to proving a lower bound on $\mathbf{P}\{M_n \leq m\}$, namely, bounding $\mathbf{P}\{v_n \in N_{n,m}\}$, then applying a union bound, will not work. It is the approach we used in Section 1.1 when discussing many *independent* random walks, and only begins to yield tail bounds when $m \leq m^* + O(1)$.

On the other hand, we observe that we can *easily* prove strong enough bounds for one group of potential nodes of $N_{n,m}$ when $m \leq m' + O(1)$, namely, the set $G_{n,m}$ (we recall that $m' = \Lambda'(t)n - (3 \log n)/2t$). By Lemma 11 (or, indeed, by Lemma 7), when m is not too far from m' , $\mathbf{P}\{v_h \in G_{n,m}\} = O(\mathbf{P}\{v_h \in N_{n,m}\}/n)$, from which strong bounds on $\mathbf{P}\{G_{n,m} \neq \emptyset\}$ follow directly from Theorem 1 and a union bound. Intuitively, this bound on $G_{n,m}$ contains almost the whole “reason for” the lower bound, in the following sense. Given a node v at depth n and $m \leq m'$, if $S_v \leq m$ is to occur, then *either* S_v is “close to being” a leading node, in the sense that there is a “small” constant a for which S_v stays above $-a$, or S_v is “far from being” a leading node, in which case there is a “large” constant c for which S_v does *not* stay above $-c$ (or something in between these two scenarios occurs). The former scenario is unlikely due to the bounds in Lemma 11, and the latter scenario is unlikely due to large deviations estimates (and everything in between is unlikely due to a mix of these two reasons).

We prove our general bound by splitting $N_{n,m}$ into many groups which differentiate among the possibilities outlined in the above sketch. We then bound the probability each group is non-empty using large deviations estimates and the Lemmas from Section 4.2. When we recombine the bounds we obtain for the individual groups (by union bound), the result will be strong enough to yield bounds on $\mathbf{P}\{N_{n,m} \neq \emptyset\}$ that are exponentially small in $m' - m$ when m is not too far from m' . For convenience, we restate the lower bound we are aiming to prove.

Lemma 13. *There are $C_2 > 0$, $\delta_2 > 0$ such that for all $x > 0$ $\mathbf{P}\{M_n \leq m' - x\} \leq C_2 e^{-\delta_2 x}$.*

Theorem 3 follows easily from the above lemma and Corollary 10; its proof appears in Section 6, below. In proving Lemma 13, it will be useful to use Chernoff (1952) bounds as well as Theorem 1, as Chernoff bounds are not limited to windows of size $o(\sqrt{n})$ around a linear slope. In particular, for our choice of t we have

Lemma 14 (Chernoff Bound). *For all $r > 0$ and all integers $k \geq 1$,*

$$\mathbf{P}\{S_{v_k} \leq \Lambda'(t)k - r\} \leq \frac{e^{tr}}{d^k}.$$

Lemma 13 is a rather straightforward consequence of the following, weaker lemma, plus Lemma 14:

Lemma 15. *There are $C_3 > 0$, $\delta_3 > 0$ such that for all $m < m'$ with $m' - m \leq (2 \log n)/|t|$,*

$$\mathbf{P}\{\exists v \in N_n : m - 1 \leq S_v \leq m\} \leq C_3 e^{-\delta_3(m' - m)}. \quad (71)$$

Proof of Lemma 13 assuming Lemma 15. Given m with $0 \leq m' - m \leq \sqrt{n/g(n)}$, if $m' - m \geq 2 \log n/|t|$ then, letting $\Delta = \Lambda'(t)n - m$, by the definition of m' we have $\Delta \geq (m' - m)/4$. It follows by a union bound and by Theorem 1 that

$$\mathbf{P}\{M_n \leq m\} \leq d^n \mathbf{P}\{S_{v_n} \leq \Lambda'(t)n - \Delta\} = O(e^{t\Delta}) = O(e^{t(m' - m)/4}),$$

which proves Lemma 13 in the case that $m' - m \geq (2 \log n)/|t|$ (since t is negative). If $0 \leq m' - m < (2 \log n)/|t|$, then we have the following inclusion:

$$\{M_n \leq m\} \subseteq \left\{M_n \leq m' - \frac{2 \log n}{|t|}\right\} \cup \bigcup_{i=\lfloor m'-m \rfloor}^{\lfloor (2 \log n)/|t| \rfloor} \{\exists v \in N_n : m' - i - 1 \leq S_v \leq m' - i\}. \quad (72)$$

Since $m' - (2 \log n)/|t| = \Lambda'(t)n - (\log n)/2|t|$, by Lemma 14 and a union bound,

$$\mathbf{P} \left\{M_n \leq m' - \frac{2 \log n}{|t|}\right\} \leq \frac{1}{\sqrt{n}} = e^{-\log n/2} \leq e^{t(m'-m)/4}. \quad (73)$$

By (72), (73), Lemma 15 and a union bound, we thus have

$$\begin{aligned} \mathbf{P} \{M_n \leq m\} &\leq e^{t(m'-m)/4} + \sum_{i=\lfloor m'-m \rfloor}^{\lfloor (2 \log n)/|t| \rfloor} \mathbf{P} \{\exists v \in N_n : m' - i - 1 \leq S_v \leq m' - i\} \\ &\leq e^{t(m'-m)/4} + C_3 \sum_{i=\lfloor m'-m \rfloor}^{\lfloor (2 \log n)/|t| \rfloor} e^{-\delta_3 i} \\ &\leq e^{t(m'-m)/4} + C_3 e^{-\delta_3 \lfloor m'-m \rfloor} \left(\frac{1}{1 - e^{-\delta_3}} \right), \end{aligned}$$

which proves Lemma 13 in the case that $0 \leq m' - m \leq (2 \log n)/|t|$. \square

We now turn our attention to proving Lemma 15. We first note that we need only prove (71) for $m' - m$ larger than any fixed constant, as we may presume the bound holds for $m' - m = O(1)$ by our choice of C_3 . For the remainder of the section, we write $\Delta = \Lambda'(t)n - m$ and $\Delta' = m' - m > L$ for some large constant L . We have $\Delta' = \Delta + (3 \log n)/2|t|$, so $\Delta' \leq (2 \log n)/|t|$ and $\Delta \leq (\log n)/2|t|$.

We now proceed to define the ‘‘homogeneous’’ groups discussed above, and bound the probabilities they are non-empty, in a sequence of claims. The proof of each claim will consist of straightforward applications of Theorem 1 and Lemmas 11 and 12, and Lemma 15 will be a trivial consequence of the bounds of the claims.

Let $a = \lfloor e^{t|\Delta'/10} \rfloor$, and let A_a be the set of nodes $v \in N_n$ for which $m - 1 \leq S_v \leq m$ and for which $\{S_v \text{ abo } - a\}$ occurs.

Claim 16. $\mathbf{P} \{A_a \neq \emptyset\} = O(e^{9t(m'-m)/10})$.

Proof. By Lemma 11,

$$\mathbf{P} \{v_n \in A_a \mid m - 1 \leq S_v \leq m\} = O\left(\frac{a^9}{n}\right) = O\left(\frac{e^{9t(m'-m)/10}}{n}\right).$$

By Theorem 1, it follows that

$$\mathbf{P} \{v_n \in A_a\} = O\left(\frac{e^{9t(m'-m)/10}}{d^n}\right).$$

The claim follows by a union bound over all $v \in N_n$. \square

Next, let $c = \lfloor \log^{10} n \rfloor$, and let H_c consist of the nodes $v \in N_n$ with $m-1 \leq S_v \leq m$ and for which $\{S_v \text{ abo } - c\}$ does not occur.

Claim 17. $\mathbf{P}\{H_c \neq \emptyset\} = O\left(e^{-(m'-m)}\right)$.

Proof. If $H_c \neq \emptyset$ then for some $0 < k < n$ there is a node $y \in N_k$ for which $S_y \leq mk/n - c$. We have

$$\frac{mk}{n} - c = \frac{(\Lambda'(t)n - \Delta)k}{n} - c = \Lambda'(t)k - \frac{\Delta k}{n} - c.$$

Therefore, by Lemma 14,

$$\mathbf{P}\left\{S_{v_k} \leq \frac{mk}{n} - c\right\} = O\left(\frac{e^{t(c+\Delta k/n)}}{d^k}\right). \quad (74)$$

Since $c = \lfloor \log^{10} n \rfloor$, we may assume n is large enough that $c - \Delta k/n \geq \log^9 n$, so (74) yields

$$\mathbf{P}\left\{S_{v_k} \leq \frac{mk}{n} - c\right\} = O\left(\frac{e^{t \log^9 n}}{d^k}\right). \quad (75)$$

By a union bound, we obtain

$$\begin{aligned} \mathbf{P}\{H_c \neq \emptyset\} &\leq \sum_{k=1}^{n-1} d^k \mathbf{P}\left\{S_{v_k} \leq \frac{mk}{n} - c\right\} \\ &= O\left(\sum_{k=1}^{n-1} e^{t \log^9 n}\right) = O(e^{\log n + t \log^9 n}) = O(e^{-(m'-m)}) \end{aligned}$$

□

For each integer b with $a \leq b \leq c$, we let M_b be the set of vertices $v \in N_n$ for which $m-1 \leq S_v \leq m$ and for which $\{S_v \text{ abo } - (b+1)\}$ occurs but $\{S_v \text{ abo } - (b+1)\}$ does not occur. We next define a subset of M_b which we call M_b^{mid} by saying that $v_n \in M_b^{\text{mid}}$ if $v_n \in M_b$ and additionally, there is k with $b^{57} \leq k \leq n - b^{57}$ for which $S_{v_k} \leq mk/n - b$. We extend this definition to all nodes $v \in N_n$ by symmetry.

Similarly, for values k with $k < b^{57}$ or with $n - k < b^{57}$, we define a set M_b^k by saying that $v_n \in M_b^k$ if $v_n \in M_b$ and additionally, k is the *smallest value* for which $S_{v_k} \leq mk/n - b$; again, we extend this definition to all nodes $v \in N_n$ by symmetry. The sets M_b^{mid} and $\{M_b^k : \min(k, n-k) < b^{57}\}$ partition M_b . We now bound the probabilities that these sets are non-empty.

Claim 18. $\mathbf{P}\left\{\bigcup_{b=a}^c \{M_b^{\text{mid}} \neq \emptyset\}\right\} = O(e^{t(m'-m)/11})$.

Proof. Fix an integer b with $a \leq b \leq c$. By Lemma 12,

$$\mathbf{P}\left\{v_n \in M_b^{\text{mid}} \mid m-1 \leq S_{v_n} \leq m\right\} = O\left(\frac{1}{nb^9}\right),$$

so by Theorem 1,

$$\mathbf{P}\left\{v_n \in M_b^{\text{mid}}\right\} = O\left(\frac{\mathbf{P}\{m-1 \leq S_{v_n} \leq m\}}{nb^9}\right) = O\left(\frac{ne^{t(m'-m)}}{nb^9 d^n}\right) = O\left(\frac{e^{t(m'-m)}}{b^9 d^n}\right).$$

By a union bound, it follows that

$$\mathbf{P} \left\{ M_b^{\text{mid}} \neq \emptyset \right\} = O \left(\frac{e^{t(m'-m)}}{b^9} \right),$$

so by summing over b with $a \leq b \leq c$, since $a = e^{\lfloor t(m'-m)/11 \rfloor}$ we obtain

$$\mathbf{P} \left\{ \bigcup_{b=a}^c \{M_b^{\text{mid}} \neq \emptyset\} \right\} = O \left(\sum_{b=a}^c \frac{e^{t(m'-m)}}{b^9} \right) = O \left(\frac{e^{t(m'-m)}}{a^{10}} \right) = O(e^{t(m'-m)/11}).$$

□

Claim 19. $\mathbf{P} \left\{ \bigcup_{b=a}^c \bigcup_{k=1}^{b^{57}-1} \{M_b^k \neq \emptyset\} \right\} = O(e^{-(m'-m)})$.

Proof. Fix b and k as above. For each node x at depth k , let W_x be the set of descendants of x in M_b^k . By a union bound over N_k , $\mathbf{P} \{M_b^k \neq \emptyset\} \leq d^k \mathbf{P} \{W_{v_k} \neq \emptyset\}$. If W_{v_k} is nonempty, then necessarily $mk/n - (b+1) \leq S_{v_k} \leq mk/n - b$. Since $k \leq b^{57} \leq c^{57} \leq (\log n)^{171}$,

$$\frac{mk}{n} - b = \Lambda'(t)k + \frac{\Delta k}{n} - b = \Lambda'(t)k - b + o(1).$$

It follows by Lemma 14 that

$$\begin{aligned} \mathbf{P} \{W_{v_k} \neq \emptyset\} &= \Theta(\mathbf{P} \{S_{v_k} \leq \Lambda'(t)k - b\}) \\ &= O \left(\frac{e^{tb}}{d^k} \right). \end{aligned} \tag{76}$$

By (76) and a union bound, we have that $\mathbf{P} \{M_b^k \neq \emptyset\} = O(e^{tb})$, so by a second union bound

$$\mathbf{P} \left\{ \bigcup_{b=a}^c \bigcup_{k=1}^{b^{57}-1} \{M_b^k \neq \emptyset\} \right\} = O \left(\sum_{b=a}^c b^{57} e^{tb} \right) = O(a^{57} e^{ta}) = O(e^{-(m'-m)}).$$

□

Claim 20. $\mathbf{P} \left\{ \bigcup_{b=a}^c \bigcup_{k=1}^{b^{57}-1} \{M_b^{n-k} \neq \emptyset\} \right\} = O(e^{-(m'-m)})$.

Proof. Fix b and k as above. For each node x at depth $n-k$, let W_x be the set of descendants of x in M_b^{n-k} . By a union bound,

$$\mathbf{P} \left\{ M_b^{n-k} \neq \emptyset \right\} \leq d^{n-k} \mathbf{P} \{W_{v_{n-k}} \neq \emptyset\}. \tag{77}$$

Suppose W_{v_r} is nonempty – then necessarily

$$\frac{m(n-k)}{n} - (b+1) \leq S_{v_{n-k}} \leq \frac{m(n-k)}{n} - b,$$

and in addition $\{S_{v_{n-k}} \text{ abo } - b\}$ must occur (or else $n - k$ is not the *first* time the random walk ending at v_n falls b below its conditional mean). Now, by Theorem 1, since $\Delta \geq (3 \log n)/2t$,

$$\begin{aligned} \mathbf{P} \left\{ S_{v_{n-k}} \leq \frac{m(n-k)}{n} - b \right\} &= \mathbf{P} \left\{ S_{v_{n-k}} \leq \Lambda'(t)(n-k) - \frac{\Delta(n-k)}{n} - b \right\} \\ &= \Theta \left(\frac{e^{t(\Delta(n-k)/n+b)}}{d^{n-k} \sqrt{n-k}} \right) = \Theta \left(\frac{e^{t(\Delta+b)}}{d^{n-k} \sqrt{n}} \right) = O \left(\frac{ne^{tb}}{d^{n-k}} \right). \end{aligned} \tag{78}$$

Furthermore, by Lemma 11,

$$\mathbf{P} \left\{ S_{v_{n-k}} \text{ abo } - b \mid \frac{m(n-k)}{n} - (b+1) \leq S_{v_{n-k}} \leq \frac{m(n-k)}{n} - b \right\} = O \left(\frac{b^9}{n-k} \right) = O \left(\frac{b^9}{n} \right). \tag{79}$$

Combining (78) and (79) yields that

$$\mathbf{P} \{W_{v_{n-k}} \neq \emptyset\} = O \left(\frac{b^9 e^{tb}}{d^{n-k}} \right) = O \left(\frac{e^{9 \log b + tb}}{d^{n-k}} \right),$$

which combined with (77) implies that $\mathbf{P} \{M_b^{n-k} \neq \emptyset\} = O(e^{9 \log b + tb})$. Just as in the proof of Claim 19, summing this bound over b and k yields the result. \square

We are now prepared for

Proof of Lemma 15. It is immediate from the definitions of the sets A_a , H_c , M_b^{mid} and M_b^k that any vertex $v \in N_n$ with $m-1 \leq S_v \leq m$ is either in A_a , or in H_c , or in M_b^{mid} (for some integer $a \leq b \leq c$), or in M_b^k (for some integer $a \leq b \leq c$ and some integer k for which either $1 \leq k \leq b^{33}$ or $h - b^{33} \leq k \leq h$). Applying Claims 16, 17, 18, 19, and 20, respectively, to bound each of these events, it follows that

$$\begin{aligned} \mathbf{P} \{\exists v \in N_n : m-1 \leq S_v \leq m\} &= O(e^{9t(m'-m)/10}) + O(e^{-(m'-m)}) + O(e^{t(m'-m)/11}) \\ &\quad + O(e^{-(m'-m)}) + O(e^{-(m'-m)}) \\ &= O(e^{-\min(|t|/11, 1)(m'-m)}), \end{aligned}$$

so the bound of Lemma 15 holds as long as we choose δ_1 so that $0 < \delta_1 \leq \min(|t|/11, 1)$ and choose C_1 large enough. \square

6 Proof of Theorem 3

By Corollary 10, there exist $C_1 > 0$, $\delta_1 > 0$ such that for all $x > 0$,

$$\mathbf{P} \{M_n \geq m' + x \mid \mathcal{S}\} \leq C_1 e^{-\delta_1 x}.$$

By Lemma 13 and Bayes' formula, there exist $C_2 > 0$, $\delta_2 > 0$ such that for all $x > 0$,

$$\mathbf{P} \{M_n \leq m' - x \mid \mathcal{S}\} \leq \frac{\mathbf{P} \{M_n \leq m' - x\}}{\mathbf{P} \{\mathcal{S}\}} \leq \frac{C_2}{\mathbf{P} \{\mathcal{S}\}} e^{-\delta_2 x}.$$

Taking $C_4 = \max\{C_1, C_2/\mathbf{P} \{\mathcal{S}\}\}$ and $\delta = \min\{\delta_1, \delta_2\}$, we obtain that for all $x > 0$,

$$\mathbf{P} \{|M_n - m'| \geq x \mid \mathcal{S}\} \leq C_4 e^{-\delta x}.$$

It follows immediately that

$$|\mathbf{E}\{M_n \mid \mathcal{S}\} - m'| \leq \mathbf{E}\{|M_n - m'| \mid \mathcal{S}\} \leq \sum_{i=0}^{\infty} \mathbf{P}\{|M_n - m'| \geq i \mid \mathcal{S}\} \leq \frac{C_4}{1 - e^{-\delta}},$$

proving (6); and letting $\Delta = C_4/(1 - e^{-\delta})$, for all $x > 0$ we have

$$\mathbf{P}\{|M_n - \mathbf{E}\{M_n \mid \mathcal{S}\}| > x \mid \mathcal{S}\} \leq \mathbf{P}\{|M_n - m'| > x - \Delta \mid \mathcal{S}\} \leq C_4 e^{\Delta\delta} e^{-\delta x},$$

so (7) holds with this choice of δ and with $C = C_4 e^{\Delta\delta}$.

7 Acknowledgements

The authors would like to thank an anonymous referee for many useful comments and suggestions.

References

- Noga Alon and Joel H. Spencer. *The Probabilistic Method*. John Wiley & Sons, Inc, second edition, 2000.
- Erik Sparre Andersen. Fluctuations of sums of random variables. *Mathematica Scandinavica*, 1:263–285, 1953.
- Markus Bachmann. Limit theorems for the minimal position in a branching random walk with independent logconcave displacements. *Advances in Applied Probability*, 32(1):159–176, 2000.
- Markus Bachmann. *Limit theorems for the minimal position in a branching random walk with independent logconcave displacements*. PhD thesis, Purdue University, 1998.
- R.R. Badahur and R. Ranga Rao. On deviations of the sample mean. *Annals of Mathematical Statistics*, 31(4):1015–1027, December 1960.
- J. D. Biggins. The first- and last-birth problems for a multitype age-dependent branching process. *Advances in Applied Probability*, 8(3):446–459, 1976.
- Maury Bramson. Maximum displacement of branching Brownian motion. *Communications on Pure and Applied Mathematics*, 31:531–581, 1978a.
- Maury Bramson. Minimal displacement of branching random walk. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 55:89–108, 1978b.
- Maury Bramson and Ofer Zeitouni. Tightness for a family of recursive equations. submitted, 2006.
- Maury Bramson and Ofer Zeitouni. Tightness for the minimal displacement of branching random walk. *Journal of Statistical Mechanics: Theory and Experiment*, July 2007, 2007. URL http://www.iop.org/EJ/article/1742-5468/2007/07/P07010/jstat7_07_p07010.html.

- N. Broutin and L. Devroye. Large deviations for the weighted height of an extended class of trees. *Algorithmica*, 46:271–279, 2006.
- N. Broutin, L. Devroye, and E. McLeish. Weighed height of random trees. *Acta Informatica*, 45(237-277), 2008.
- B. Chauvin and M. Drmota. The random multisection problem, travelling waves, and the distribution of the height of m-ary search trees. *Algorithmica*, 46:299–327, 2006.
- H. Chernoff. A measure of the asymptotic efficiency for tests of a hypothesis based on the sum of observables. *Ann. Math. Statist.*, 2:493–509, 1952.
- K.L. Chung and P. Erdős. On the application of the Borel-Cantelli lemma. *Transactions of the American Mathematical Society*, 72(1), 1952.
- F.M. Dekking and B. Host. Limit distributions for minimal displacement of branching random walks. *Zeitschrift für Wahrscheinlichkeitstheorie und verwandte Gebiete*, 90(3):403–426, 1991.
- A. Dembo and O. Zeitouni. *Large Deviation Techniques and Applications*. Jones and Bartlett, 1992.
- L. Devroye. Branching processes and their application in the analysis of tree structures and tree algorithms. In M. Habib, C. McDiarmid, J. Ramirez-Alfonsin, and B. Reed, editors, *Probabilistic Methods for Algorithmic Discrete Mathematics*, volume 16 of *Springer Series on Algorithms and Combinatorics*, pages 249–314, Berlin, 1998. Springer.
- L. Devroye and B. Reed. On the variance of the height of random binary search trees. *SIAM Journal on Computing*, 24:1157–1162, 1995.
- M. Drmota. An analytic approach to the height of binary search trees II. *Journal of the ACM*, 50:333–374, 2003.
- J.M. Hammersley. Postulates for subadditive processes. *The Annals of Probability*, 2(4):652–680, 1974.
- T. E. Harris. *The Theory of Branching Processes*. Springer, Berlin, 1963.
- Yueyun Hu and Zhan Shi. Minimal position and critical martingale convergence in branching random walks, and directed polymers on disordered trees. Submitted, February 2007.
- J.F.C. Kingman. The first birth problem for an age-dependent branching process. *The Annals of Probability*, 3(5):790–801, 1975.
- M.A. Lifshits. Some limit theorems on binary trees. in preparation, 2007.
- C. McDiarmid. Minimal positions in a branching random walk. *Advances in Applied Probability*, 5(1):128–139, 1995.
- B. Reed. The height of a random binary search tree. *Journal of the ACM*, 50:306–332, 2003.