

THE MAXIMUM NUMBER OF CLIQUES IN A GRAPH EMBEDDED IN A SURFACE

VIDA DUJMOVIĆ, GAŠPER FIJAŽ, GWENAËL JORET, AND DAVID R. WOOD

ABSTRACT. This paper studies the following question: Given a surface Σ and an integer n , what is the maximum number of cliques in an n -vertex graph embeddable in Σ ? We characterise the extremal graphs for this question, and prove that the answer is between $8(n - \omega) + 2^\omega$ and $8n + \frac{3}{2} 2^\omega + o(2^\omega)$, where ω is the maximum integer such that the complete graph K_ω embeds in Σ . For the surfaces $\mathbb{S}_0, \mathbb{S}_1, \mathbb{S}_2, \mathbb{N}_1, \mathbb{N}_2, \mathbb{N}_3$ and \mathbb{N}_4 we establish an exact answer.

MSC Classification: 05C10 (Topological graph theory), 05C35 (Extremal problems)

1. INTRODUCTION

A *clique* in a graph¹ is a set of pairwise adjacent vertices. Let $c(G)$ be the number of cliques in a graph G . This paper studies the following question at the intersection of topological and extremal graph theory: Given a surface Σ and an integer n , what is the maximum number of cliques in an n -vertex graph embeddable in Σ ?

For previous bounds on the maximum number of cliques in certain graph families see [4, 10, 17, 18] for example. For background on graphs embedded in surfaces see [8, 16]. Every surface is homeomorphic to \mathbb{S}_g , the orientable surface with g handles, or to \mathbb{N}_h , the non-orientable surface with h crosscaps. The *Euler characteristic* of \mathbb{S}_g is $2 - 2g$. The *Euler characteristic* of \mathbb{N}_h is $2 - h$. Throughout the paper, fix a surface Σ with Euler characteristic χ . If $\Sigma = \mathbb{S}_0$ then let $\omega = 3$, otherwise let ω be the maximum integer such that the complete graph K_ω embeds in Σ . To avoid trivial exceptions, we implicitly assume that $|V(G)| \geq 3$ whenever $\Sigma = \mathbb{S}_0$.

Our first main result is to characterise the n -vertex graphs embeddable in Σ with the maximum number of cliques; see Theorem 2 in Section 2. Using this result we determine an exact formula for the maximum number of cliques in an n -vertex graph embeddable in each of the the sphere \mathbb{S}_0 , the torus \mathbb{S}_1 , the double torus \mathbb{S}_2 , the projective plane \mathbb{N}_1 , the Klein bottle \mathbb{N}_2 , as well as \mathbb{N}_3 and \mathbb{N}_4 ; see Section 3. Our third main result estimates the maximum number

Date: June 22, 2009.

Vida Dujmović is supported by the Natural Sciences and Engineering Research Council of Canada. Gašper Fijaž is supported in part by the Slovenian Research Agency, Research Program P1-0297. Gwenaël Joret is a Postdoctoral Researcher of the Fonds National de la Recherche Scientifique (F.R.S.-FNRS). David Wood is supported by a QEII Research Fellowship from the Australian Research Council.

¹We consider simple, finite, undirected graphs G with vertex set $V(G)$ and edge set $E(G)$.

of cliques in terms of ω . We prove that the maximum number of cliques in an n -vertex graph embeddable in Σ is between $8(n - \omega) + 2^\omega$ and $8n + \frac{3}{2}2^\omega + o(2^\omega)$; see Theorem 13 in Section 4.

2. CHARACTERISATION OF EXTREMAL GRAPHS

The upper bounds proved in this paper are of the form: every graph G embeddable in Σ satisfies $c(G) \leq 8|V(G)| + f(\Sigma)$ for some function f . Define the *excess* of G to be $c(G) - 8|V(G)|$. Thus the excess of G is at most Q if and only if $c(G) \leq 8|V(G)| + Q$. Theorem 13 proves that the maximum excess of a graph embeddable in Σ is finite.

In this section we characterise the graphs embeddable in Σ with maximum excess. A *triangulation* of Σ is an embedding of a graph in Σ in which each face is bounded by a triangle. (We assume that every face of a graph embedding is homeomorphic to a unit disc.)

Lemma 1. *Every graph G embeddable in Σ with maximum excess is a triangulation of Σ .*

Proof. Since adding edges increases the number of cliques, the vertices on the boundary of each face of G form a clique.

Suppose that some face f of G has at least four distinct vertices in its boundary. Let G' be the graph obtained from G by adding one new vertex adjacent to four distinct vertices of f . Thus G' is embeddable in Σ , has $|V(G)| + 1$ vertices, and has $c(G) + 16$ cliques, which contradicts the choice of G . Now assume that every face of G has at most three distinct vertices.

Suppose that some face f of G is not a triangle. Thus the facial walk of f contains vertices u, v, w, v in this order (where v is repeated in f). Let G' be the graph obtained from G by adding two new vertices p and q , where p is adjacent to $\{u, v, w, q\}$, and q is adjacent to $\{u, v, w, p\}$. So G' is embeddable in Σ and has $|V(G)| + 2$ vertices. If $S \subseteq \{p, q\}$ and $S \neq \emptyset$ and $T \subseteq \{u, v, w\}$, then $S \cup T$ is a clique of G' but not of G . It follows that G' has $c(G) + 24$ cliques, which contradicts the choice of G . Hence G is a triangulation of Σ . □

Let G be a triangulation of Σ . An edge of G is *reducible* if it is in exactly two triangles of G . We say G is *irreducible* if no edge of G is reducible [2, 3, 5, 6, 9, 12, 14, 15]. Note that K_3 is a triangulation of \mathbb{S}_0 , and by the above definition, K_3 is irreducible. In fact, it is the only irreducible triangulation of \mathbb{S}_0 . We take this somewhat non-standard approach so that Theorem 2 holds for all surfaces.

Let vw be a reducible edge of G . Let vwx and $vw y$ be the two faces incident to vw in G . As illustrated in Figure 1, let G/vw be the graph obtained from G by *contracting* vw ; that is, delete the edges vw, wy, wx , and identify v and w into v . G/vw is a simple graph since x and y are the only common neighbours of v and w . Indeed, G/vw is a triangulation of Σ .

Conversely, we say that G is obtained from G/vw by *splitting* the path xvy at v . If, in addition, $xy \in E(G)$, then we say that G is obtained from G/vw by *splitting* the triangle xvy at v . Note that splitting a face xvy simply means to add a new vertex adjacent to each of x, v, y .




FIGURE 1. Contracting a reducible edge.

Graphs embeddable in Σ with maximum excess are characterised in terms of irreducible triangulations as follows.

Theorem 2. *Let X be the set of irreducible triangulations of Σ with maximum excess Q . Then the excess of every graph G embeddable in Σ is at most Q , with equality if and only if G is obtained from some graph in X by repeatedly splitting triangles.*

Proof. We proceed by induction on $|V(G)|$. By Lemma 1, we may assume that G is a triangulation of Σ . If G is irreducible, then the claim follows from the definition of X and Q . Otherwise, some edge vw of G is in exactly two triangles vwx and $vw y$. By induction, the excess of G/vw is at most Q , with equality if and only if G/vw is obtained from some $H \in X$ by repeatedly splitting triangles. Hence $c(G/vw) \leq 8|V(G/vw)| + Q$.

Observe that every clique of G that is not in G/vw is in $\{A \cup \{w\} : A \subseteq \{x, v, y\}\}$. Thus $c(G) \leq c(G/vw) + 8$, with equality if and only if xvy is a triangle. Hence $c(G) \leq 8|V(G)| + Q$; that is, the excess of G is at most Q .

Now suppose that the excess of G equals Q . Then the excess of G/vw equals Q , and $c(G) = c(G/vw) + 8$ (implying xvy is a triangle). By induction, G/vw is obtained from H by repeatedly splitting triangles. Therefore G is obtained from H by repeatedly splitting triangles.

Conversely, suppose that G is obtained from some $H \in X$ by repeatedly splitting triangles. Then xvy is a triangle and G/vw is obtained from H by repeatedly splitting triangles. By induction, the excess of G/vw equals Q , implying the excess of G equals Q . \square

3. LOW-GENUS SURFACES

To prove an upper bound on the number of cliques in a graph embedded in Σ , by Theorem 2, it suffices to consider irreducible triangulations of Σ with maximum excess. The complete list of irreducible triangulations is known for $\mathbb{S}_0, \mathbb{S}_1, \mathbb{S}_2, \mathbb{N}_1, \mathbb{N}_2, \mathbb{N}_3$ and \mathbb{N}_4 . In particular, Steinitz and

TABLE 1. The maximum excess of an n -vertex irreducible triangulation of Σ .

Σ	χ	ω	$n = 3$	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	max	
\mathbb{S}_0	2	3	-16																			-16
\mathbb{S}_1	0	7			72	48	40	32														72
\mathbb{S}_2	-2	8						208	160	136	128	120	96	88	80							208
\mathbb{N}_1	1	6		16	8																	16
\mathbb{N}_2	0	6				48	48	40	32													48
\mathbb{N}_3	-1	7					104	104	96	80	80	72	64	56								104
\mathbb{N}_4	-2	8					216	208	152	136	136	136	128	120	112	107	99	91	83	75		216

Rademacher [11] proved that K_3 is the only irreducible triangulation of \mathbb{S}_0 (under our definition of irreducible). Lavrenchenko [5] proved that there are 21 irreducible triangulations of \mathbb{S}_1 , each with between 7 and 10 vertices; see [7] for related results. Sulanke [12] proved that there are 396,784 irreducible triangulations of \mathbb{S}_2 , each with between 10 and 17 vertices. Barnette [1] proved that the embeddings of K_6 and $K_7 - K_3$ in \mathbb{N}_1 are the only irreducible triangulations of \mathbb{N}_1 . Sulanke [15] proved that there are 29 irreducible triangulations of \mathbb{N}_2 , each with between 8 and 11 vertices (correcting an earlier result by Lawrencenko and Negami [6]). Sulanke [12] proved that there are 9708 irreducible triangulations of \mathbb{N}_3 , each with between 9 and 16 vertices. Sulanke [12] proved that there are 6,297,982 irreducible triangulations of \mathbb{N}_4 , each with between 9 and 22 vertices. Using the lists of all irreducible triangulations due to Sulanke [13] and a naive algorithm for counting cliques², we have computed the set X in Theorem 2 for each of the above surfaces; see Table 1. This data with Theorem 2 implies the following results.

Proposition 3. Every planar graph G with $|V(G)| \geq 3$ has at most $8|V(G)| - 16$ cliques, as proved by Wood [17]. Moreover, a planar graph G has $8|V(G)| - 16$ cliques if and only if G is obtained from the embedding of K_3 in \mathbb{S}_0 by splitting triangles.

Proposition 4. Every toroidal graph G has at most $8|V(G)| + 72$ cliques. Moreover, a toroidal graph G has $8|V(G)| + 72$ cliques if and only if G is obtained from the embedding of K_7 in the torus by splitting triangles.

Proposition 5. Every graph G embeddable in \mathbb{S}_2 has at most $8|V(G)| + 208$ cliques. Moreover, a graph G embeddable in \mathbb{S}_2 has $8|V(G)| + 208$ cliques if and only if G is obtained from one of the following two graph embeddings by splitting triangles³:

²The code is available from the authors upon request.

³This representation describes a graph with vertex set $\{a, b, c, \dots\}$ by adjacency lists of the vertices in order a, b, c, \dots . The graph # refers to the position in Sulanke's file [13].

graph #1: bcde,afdghic,abiehfgd,acgbfihe,adhcgfb,begchjid,bdcfeijh,bgjfcedi,bhdfjgec,fhgi
graph #6: bcde,afdghijc,abjehfgd,acgbfjihe,adhcgjfb,begchjd,bdcfejh,bgjfcedi,bhdj,bidfhgec

Proposition 6. Every projective planar graph G has at most $8|V(G)| + 16$ cliques. Moreover, a projective planar graph G has $8|V(G)| + 16$ cliques if and only if G is obtained from the embedding of K_6 in \mathbb{N}_1 by splitting triangles.

Proposition 7. Every graph G embeddable in \mathbb{N}_2 has at most $8|V(G)| + 48$ cliques. Moreover, a graph G embeddable in \mathbb{N}_2 has $8|V(G)| + 48$ cliques if and only if G is obtained from one of the following three graph embeddings by splitting triangles:

graph #3: bcdef,afgdhec,abefd,acfhbge,adghbcf,aecdhgb,bfhed,bdfge
graph #6: bcde,afdghc,abhegd,acgbfhe,adhcgfb,beghd,bdcefh,bgfdec
graph #26: bcdef,afghidec,abefd,acfhgibe,adbcf,aecdhighb,bfidh,bgdfi,bhfgd

Proposition 8. Every graph G embeddable in \mathbb{N}_3 has at most $8|V(G)| + 104$ cliques. Moreover, a graph G embeddable in \mathbb{N}_3 has $8|V(G)| + 104$ cliques if and only if G is obtained from one of the following 15 graph embeddings by splitting triangles:

graph #1: bcde,afdghic,abiegfd,acfbgie,adicghfb,behigcd,bdifceh,bgefi,bhfgdec
graph #3: bcde,afdghic,abiehd,achfbgie,adichgfb,begihd,bdifeh,bgecdfi,bhfgdec
graph #4: bcde,afdghic,abiehd,achifbge,adgichfb,behgid,bdeifh,bgfecdi,bhdfgec
graph #6: bcde,afdghic,abiehfd,acfbgihe,adhcgfb,beighcd,bdifh,bgfecdi,bhdfgec
graph #8: bcde,afdghic,abiehfgd,acfbgihe,adhcgfb,begcd,bdiefch,bgcedi,bhdgec
graph #10: bcde,afdghic,abifegd,acgbfhie,adigcfb,becighd,bdceifh,bgfdi,bhdegfc
graph #12: bcde,afdghic,abifehd,achfbgie,adihcfb,becighd,bdifh,bgfdcei,bhedgfc
graph #14: bcde,afdghic,abigehd,achfbgie,adihcgfb,beghd,bdicefh,bgfdcei,bhedgfc
graph #16: bcde,afdghic,abiegd,acgbhfe,adicghfb,behdig,bfihecd,bdfegi,bhgfdec
graph #19: bcde,afghdic,abiehd,achbifge,adgichfb,behidg,bfdeih,bgifecd,bdfhgec
graph #20: bcde,afghdic,abigehd,achbifge,adgichfb,beidg,bfdecih,bgiecd,bdfeghc
graph #21: bcde,afghdic,abihegd,acgfbhie,adigchfb,behdg,bfdceih,bgicefd,bdeghe
graph #22: bcde,afghdic,abihegd,acgfbhie,adhcgfb,beidg,bfdceih,bgiced,bdfeghc
graph #82: bcdef,afgdhiec,abefd,acfigbhe,adhgibcf,aecdihgb,bfheid,bdegfi,bhfdge
graph #2464: bcdef,afghijdec,abefd,acfhigjbe,adbcf,aecdhighb,bfidjh,bgjfdi,bhdfgj,bifhgd

Proposition 9. Every graph G embeddable in \mathbb{N}_4 has at most $8|V(G)| + 216$ cliques. Moreover, a graph G embeddable in \mathbb{N}_4 has $8|V(G)| + 216$ cliques if and only if G is obtained from one of the following three graph embeddings by repeatedly splitting triangles:

graph #1: bcdef,afdgehic,abiegfhd,achgbfie,adicgbhf,aehcgidb,bdhifce,befcdgi,bhgfdec
graph #2: bcdef,afdgehic,abifehgd,acgbfhie,adigbhcf,aecighdb,bdchfie,becgfdi,bhdegfc

4. A BOUND FOR ALL SURFACES

We start with the following upper bound on the minimum degree of a graph.

Lemma 10. *Every graph G embedded in Σ has minimum degree at most*

$$6 + \frac{\omega^2 - 7\omega + 10}{|V(G)|} .$$

Proof. If $\Sigma = \mathbb{S}_g$, then $g = \frac{2-\chi}{2} = \lceil \frac{(\omega-3)(\omega-4)}{12} \rceil$. If $\Sigma = \mathbb{N}_h$, then $h = 2 - \chi = \lceil \frac{(\omega-3)(\omega-4)}{6} \rceil$, unless $h = 3$ in which case $\omega = 7$. Since $(\omega - 3)(\omega - 4)$ is even, it follows that⁴

$$(1) \quad -6\chi \leq \omega^2 - 7\omega + 10 .$$

Say G has minimum degree d . It follows from Euler's Formula that $|E(G)| \leq 3|V(G)| - 3\chi$. By (1),

$$d \leq \frac{2|E(G)|}{|V(G)|} \leq \frac{6|V(G)| - 6\chi}{|V(G)|} \leq 6 + \frac{\omega^2 - 7\omega + 10}{|V(G)|} . \quad \square$$

For graphs in which the number of vertices is slightly more than ω , Lemma 10 can be reinterpreted as follows.

Lemma 11. *Assume $\omega \geq 18$. Let $s := \lceil \sqrt{\omega - 2} \rceil - 4 \geq 0$ and let $j \in [0, s]$. Then every graph G embeddable in Σ with at least $\omega + j$ vertices has minimum degree at most $\omega - j - 1$.*

Proof. Say G has minimum degree d . By Lemma 10,

$$d \leq 6 + \frac{\omega^2 - 7\omega + 10}{\omega + j} = \omega - j - 1 + \frac{10 + j(j + 7)}{\omega + j} .$$

The assumption that $j \leq s \leq \lceil \sqrt{\omega - 2} \rceil - 4$ implies that $j + 3 < \sqrt{\omega - 2}$ and $j^2 + 6j + 10 < \omega$. That is, $10 + j(j + 7) < \omega + j$. Hence $d \leq \omega - j - 1$. \square

Now we prove our first upper bound on the number of cliques.

Lemma 12. *Assume $\omega \geq 18$. Let $s := \lceil \sqrt{\omega - 2} \rceil - 4 \geq 0$. Let G be an n -vertex graph embeddable in Σ . Then*

$$c(G) \leq 2^\omega + 2^{\omega-1} + (n - \omega - s)2^{\omega-s-1} .$$

Proof. Let v_1, v_2, \dots, v_n be an ordering of the vertices of G such that v_i has minimum degree in the subgraph $G_i := G - \{v_1, \dots, v_{i-1}\}$. Let d_i be the degree of v_i in G_i (which equals the minimum degree of G_i). Charge each clique C in G to the vertex $v_i \in C$ with i minimum.

⁴Note that (1) holds with equality if $\Sigma = \mathbb{S}_g$ and $g = 12k^2 - 3k + 1$ and $\omega = 12k + 2$ for some $k \in \mathbb{Z}^+$.

We distinguish three types of vertices. Vertex v_i is type-1 if $i \in [1, n - \omega - s]$. Vertex v_i is type-2 if $i \in [n - \omega - s + 1, n - \omega]$. Vertex v_i is type-3 if $i \in [n - \omega + 1, n]$. Note that if $s = 0$ then there are no type-2 vertices.

Each clique charged to a type-3 vertex is contained in $\{v_{n-\omega+1}, \dots, v_n\}$, and there are at most 2^ω such cliques.

Say C is a clique charged to a type-1 or type-2 vertex v_i . Then $C - \{v_i\}$ is contained in $N_{G_i}(v_i)$, which consists of d_i vertices. Thus the number of cliques charged to v_i is at most 2^{d_i} . Recall that d_i equals the minimum degree of G_i , which has $n - i + 1$ vertices.

If v_i is type-2 then $d_i \leq \omega - j - 1$ by Lemma 11 with $j = n - \omega - i + 1 \in [1, s]$. Thus the number of cliques charged to type-2 vertices is at most

$$\sum_{j=1}^s 2^{\omega-j-1} \leq \sum_{j=1}^{\omega-2} 2^j < 2^{\omega-1} .$$

If v_i is type-1 then G_i has more than $\omega + s$ vertices, and thus $d_i \leq \omega - s - 1$ by Lemma 11 with $j = s$. Thus the number of cliques charged to type-1 vertices is at most $(n - \omega - s)2^{\omega-s-1}$. \square

We now prove the main result of this section.

Theorem 13. *The maximum number of cliques in an n -vertex graph embeddable in Σ is at least $8(n - \omega) + 2^\omega$ and at most $8n + \frac{3}{2}2^\omega + o(2^\omega)$.*

Proof. To prove the lower bound, start with K_ω embedded in Σ (which has 2^ω cliques), and then while there are less than n vertices, insert a new vertex adjacent to each vertex of a single face. Each new vertex adds at least 8 new cliques. Thus we obtain an n -vertex graph embedded in Σ with at least $8(n - \omega) + 2^\omega$ cliques.

To prove the upper bound, by Theorem 2, we may assume that G is an n -vertex irreducible triangulation of Σ . Nakamoto and Ota [9] proved that, in this case, $|V(G)| \leq \max\{270 - 171\chi, 4\}$. See [2] for an earlier bound, and see [3] for an improved bound in the orientable case. By (1) and since $\omega \geq 4$,

$$|V(G)| \leq \max\{270 - 171\chi, 4\} \leq \max\{270 + \frac{171}{6}(\omega^2 - 7\omega + 10), 4\} < 29\omega^2 .$$

Thus we may assume that $\omega \geq 18$. By Lemma 12,

$$c(G) \leq 2^\omega + 2^{\omega-1} + (29\omega^2 - \omega - s)2^{\omega-s-1} < \frac{3}{2}2^\omega + 29\omega^2 2^{\omega-s-1} < \frac{3}{2}2^\omega + 2^{\omega-s+2\log\omega+4} .$$

We have $c(G) \leq \frac{3}{2}2^\omega + o(2^\omega)$ since $s \in \Theta(\sqrt{\omega})$. \square

5. CONCLUDING CONJECTURES

We conjecture that the upper bound in Theorem 13 can be improved to more closely match the lower bound.

Conjecture 14. Every graph G embeddable in Σ has at most $8|V(G)| + 2^\omega + o(2^\omega)$ cliques.

If K_ω triangulates Σ , then we conjecture the following exact answer.

Conjecture 15. Suppose that K_ω triangulates Σ . Then every graph G embeddable in Σ has at most $8(|V(G)| - \omega) + 2^\omega$ cliques, with equality if and only if G is obtained from K_ω by repeatedly splitting triangles.

By Theorem 2, this conjecture is equivalent to:

Conjecture 16. Suppose that K_ω triangulates Σ . Then K_ω is the only irreducible triangulation of Σ with maximum excess.

The results in Section 3 confirm Conjectures 15 and 16 for \mathbb{S}_0 , \mathbb{S}_1 and \mathbb{N}_1 .

Now consider surfaces possibly with no complete graph triangulation. Then the bound $c(G) \leq 8(|V(G)| - \omega) + 2^\omega$ is false for \mathbb{S}_2 , \mathbb{N}_3 and \mathbb{N}_4 . Loosely speaking, this is because these surfaces have ‘small’ ω compared to χ . In particular, except for the sphere, $\omega = \lfloor \frac{1}{2}(7 + \sqrt{49 - 24\chi}) \rfloor$, and $\omega = \frac{1}{2}(7 + \sqrt{49 - 24\chi})$ if and only if K_ω triangulates Σ . This phenomenon motivates the following conjecture:

Conjecture 17. Every graph G embeddable in Σ has at most

$$8|V(G)| - 4(7 + \sqrt{49 - 24\chi}) + 2^{(7 + \sqrt{49 - 24\chi})/2}$$

cliques, with equality if and only if K_ω triangulates Σ and G is obtained from K_ω by repeatedly splitting triangles.

There are two irreducible triangulations of \mathbb{S}_2 with maximum excess, there are three irreducible triangulations of \mathbb{N}_2 with maximum excess, there are 15 irreducible triangulations of \mathbb{N}_3 with maximum excess, and there are three irreducible triangulations of \mathbb{N}_4 with maximum excess. This suggests that for surfaces with no complete graph triangulation, a succinct characterisation of the extremal examples (as in Conjecture 16) might be difficult. Nevertheless, we conjecture the following strengthening of Conjecture 16 for all surfaces.

Conjecture 18. Every irreducible triangulation of Σ with maximum excess contains K_ω as a subgraph.

We have verified Conjectures 17 and 18 for \mathbb{S}_0 , \mathbb{S}_1 , \mathbb{S}_2 , \mathbb{N}_1 , \mathbb{N}_2 , \mathbb{N}_3 and \mathbb{N}_4 .

REFERENCES

- [1] DAVID BARNETTE. Generating the triangulations of the projective plane. *J. Combin. Theory Ser. B*, 33(3):222–230, 1982.
- [2] DAVID W. BARNETTE AND ALLAN L. EDELSON. All 2-manifolds have finitely many minimal triangulations. *Israel J. Math.*, 67(1):123–128, 1989.
- [3] SIU-WING CHENG, TAMAL K. DEY, AND SHEUNG-HUNG POON. Hierarchy of surface models and irreducible triangulations. *Comput. Geom.*, 27(2):135–150, 2004.
- [4] JÜRGEN ECKHOFF. The maximum number of triangles in a K_4 -free graph. *Discrete Math.*, 194(1–3):95–106, 1999.
- [5] SERGE A. LAVRENCENKO. Irreducible triangulations of a torus. *Ukrain. Geom. Sb.*, (30):52–62, ii, 1987. Translation in *J. Soviet Math.* 51(5):2537–2543, 1990.
- [6] SERGE LAWRENCENKO AND SEIYA NEGAMI. Irreducible triangulations of the Klein bottle. *J. Combin. Theory Ser. B*, 70(2):265–291, 1997.
- [7] JOSÉ L. MARTÍNEZ-MORALES. The irreducible triangulations of the torus as periodic tessellations of the plane. *JP J. Geom. Topol.*, 6(2):101–130, 2006.
- [8] BOJAN MOHAR AND CARSTEN THOMASSEN. *Graphs on surfaces*. Johns Hopkins University Press, Baltimore, U.S.A., 2001.
- [9] ATSUHIRO NAKAMOTO AND KATSUHIRO OTA. Note on irreducible triangulations of surfaces. *J. Graph Theory*, 20(2):227–233, 1995.
- [10] SERGUEI NORINE, PAUL SEYMOUR, ROBIN THOMAS, AND PAUL WOLLAN. Proper minor-closed families are small. *J. Combin. Theory Ser. B*, 96(5):754–757, 2006.
- [11] ERNST STEINITZ AND HANS RADEMACHER. *Vorlesungen über die Theorie der Polyeder*. Springer, Berlin, 1934.
- [12] THOM SULANKE. Generating irreducible triangulations of surfaces, 2006. <http://arxiv.org/abs/math/0606687>.
- [13] THOM SULANKE. Generating triangulations of surfaces, 2006. <http://hep.physics.indiana.edu/~tsulanke/graphs/surftri/>.
- [14] THOM SULANKE. Irreducible triangulations of low genus surfaces, 2006. <http://arxiv.org/abs/math/0606690>.
- [15] THOM SULANKE. Note on the irreducible triangulations of the Klein bottle. *J. Combin. Theory Ser. B*, 96(6):964–972, 2006.
- [16] ARTHUR T. WHITE. *Graphs, groups and surfaces*. North-Holland, 2nd edn., 1984.
- [17] DAVID R. WOOD. On the maximum number of cliques in a graph. *Graphs Combin.*, 23(3):337–352, 2007.

- [18] ALEXANDER A. ZYKOV. On some properties of linear complexes. *Mat. Sbornik N.S.*,
24(66):163–188, 1949.

SCHOOL OF COMPUTER SCIENCE

CARLETON UNIVERSITY

OTTAWA, CANADA

E-mail address: `vida@cs.mcgill.ca`

FACULTY OF COMPUTER AND INFORMATION SCIENCE

UNIVERSITY OF LJUBLJANA

LJUBLJANA, SLOVENIA

E-mail address: `gasper.fijavz@fri.uni-lj.si`

DÉPARTEMENT D'INFORMATIQUE

UNIVERSITÉ LIBRE DE BRUXELLES

BRUSSELS, BELGIUM

E-mail address: `gjoret@ulb.ac.be`

DEPARTMENT OF MATHEMATICS AND STATISTICS

THE UNIVERSITY OF MELBOURNE

MELBOURNE, AUSTRALIA

E-mail address: `woodd@unimelb.edu.au`